

Elliptic Littlewood identities

Eric M. Rains

Department of Mathematics, California Institute of Technology

June 4, 2008

Abstract

We prove analogues for elliptic interpolation functions of Macdonald's version of the Littlewood identity for (skew) Macdonald polynomials, in the process developing an interpretation of general elliptic “hypergeometric” sums as skew interpolation functions. One such analogue has an interpretation as a “vanishing integral”, generalizing a result of [9]; the structure of this analogue gives sufficient insight to enable us to conjecture elliptic versions of most of the other vanishing integrals of [9] as well. We are thus led to formulate ten conjectures, each of which can be viewed as a multivariate quadratic transformation, and can be proved in a number of special cases.

Contents

1	Introduction	1
2	Skew interpolation functions	7
3	Elliptic Cauchy identities	13
4	Elliptic Littlewood identities	18
5	Quadratic transformations	26

1 Introduction

In recent work of the author [7] (see also [1] for an independent treatment), a family of “interpolation functions” were introduced, generalizing Okounkov’s interpolation polynomials [5], which in turn generalize shifted Macdonald polynomials [12] and Macdonald polynomials [4] themselves. Among the identities satisfied by the interpolation functions is an analogue of the Cauchy identity, which for Macdonald polynomials states

$$\sum_{\mu} P_{\mu}(x_1, \dots, x_n; q, t) P_{\mu'}(y_1, \dots, y_m; q, t) = \prod_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}} (1 + x_i y_j). \quad (1.1)$$

Macdonald also proved (generalizing a result of Kadell for Jack polynomials) an analogue for Macdonald polynomials of the Littlewood identity:

$$\sum_{\mu} c_{\mu}(q, t) P_{\mu^2}(x_1, \dots, x_n; q, t) = \prod_{1 \leq i < j \leq n} \frac{(tx_i x_j; q)}{(x_i x_j; q)}, \quad (1.2)$$

where P_{λ} is a Macdonald polynomial, μ^2 denotes the partition with parts $(\mu^2)_i = \mu_{\lceil i/2 \rceil}$,

$$(x; q) := \prod_{k \geq 0} (1 - q^k x), \quad (1.3)$$

and the coefficients $c_{\mu}(q, t)$ are given by an explicit product. The purpose of the present note is to generalize this statement to the elliptic level.

The primary obstacle to such an extension is the fact that, unlike the given form of the Cauchy identity, for which the terms vanish unless the partition μ is contained in an $m \times n$ rectangle, the Littlewood identity intrinsically involves a nonterminating sum. Unfortunately, at the elliptic level, infinite sums seem inevitably to encounter convergence difficulties, making a direct extension problematical. One must thus either modify the sum in such a way as to force termination (say by a suitable choice of the coefficients c_{μ}), or replace the sum by an integral. We will, in fact, take both approaches.

Our first step is to observe that Macdonald's Littlewood identity has a generalization (implicit in [4]) to skew Macdonald polynomials, of the following form:

$$\sum_{\mu} c_{\mu}(q, t) P_{\mu^2/\lambda}(x_1, \dots, x_n; q, t) = \prod_{1 \leq i < j \leq n} \frac{(tx_i x_j; q)}{(x_i x_j; q)} \sum_{\mu} c'_{\mu}(q, t) P_{\lambda/\mu^2}(x_1, \dots, x_n; q, t). \quad (1.4)$$

Of course, this in itself makes an extension more difficult, given the absence (but see below) of a good theory of skew versions of the interpolation functions. On the other hand, the proof of Macdonald's Littlewood identity uses only the case $n = 1$ of this skew Littlewood identity, together with a corresponding case of the skew Cauchy identity. This case is particularly amenable to generalization, as both sums are finite (indeed, each has only one nonzero term), and the case $n = 1$ of skew Macdonald polynomials *does* have a very natural elliptic analogue. Indeed, the principal specialization (i.e., with variables specialized to $v, \dots, vt^{n-1}$) of a skew Macdonald polynomial can be expressed as a limit of an elliptic binomial coefficient, essentially just a value of an elliptic interpolation function. If one replaces the skew Macdonald polynomials by such elliptic binomial coefficients in the $n = 1$ case, one finds that both sums still have only one surviving term, and one is led immediately to an elliptic analogue of the identity, Lemma 4.1 below.

To obtain a more general elliptic analogue, there are two natural approaches. The first is to develop a theory of skew interpolation functions, prove a corresponding skew Cauchy identity, then directly lift Macdonald's argument to the elliptic level. Roughly speaking, skew interpolation functions should give the coefficients in a generalized branching rule:

$$\begin{aligned} \mathcal{R}_{\lambda}^{*(n+m)}(x_1, \dots, x_n, y_1, \dots, y_m; t_0, u_0; t; p, q) \\ = \sum_{\mu} \mathcal{R}_{\lambda/\mu}^{*(m)}(y_1, \dots, y_m; t_0, u_0; t; p, q) \mathcal{R}_{\mu}^{*(n)}(x_1, \dots, x_n; t_0, u_0; t; p, q). \end{aligned} \quad (1.5)$$

Since these coefficients are understood for $m = 1$, one could simply define skew interpolation functions by induction, giving an m -fold sum. However, it turns out that one can use connection coefficients together with the existence of a special case of interpolation functions expressible as a product to obtain these coefficients via a single sum. Moreover, if the arguments $y_1, \dots, y_m$ contain partial geometric progressions of step t , the coefficients of the sum simplify accordingly, and one is thus led to the definition of Section 2. A suitable analytic continuation argument gives an analogue of the Cauchy identity (Theorem 3.7), and then Macdonald's argument lifts to give an elliptic Littlewood identity, Theorem 4.4.

The other natural approach to an elliptic analogue is to retain the use of binomial coefficients (i.e., restrict one's attention to principally specialized skew Macdonald polynomials), but hope for an analogue with additional parameters. It turns out that enough degrees of freedom survive in the choice of coefficients c_μ, c'_μ that one can use those coefficients to enforce termination, giving Theorem 4.5 below. Moreover, the structure of the coefficients is such that one can analytically continue one of the two sums to a suitable contour integral, Theorem 4.6. This in turn suggests a further extension in which both sides are integrals, stated as Conjecture 2, for which we can prove a number of special cases.

The “integral=sum” version of the identity has a particularly striking interpretation coming from the fact that one can invert the elliptic binomial coefficients to move the sum inside the integral. The resulting sum of interpolation functions in the integrand then becomes a special case of the elliptic biorthogonal functions of [8, 7], and one thus deduces that a certain integral of such functions vanishes unless the indexing partition (or, rather, partition pair) has the form μ^2 . This is the elliptic analogue of a result proved for Koornwinder polynomials in [9], and in fact gives a stronger result even at the Koornwinder level, since the techniques of [9] gave no information about the nonzero values. This suggests in turn that the other results of [9] involving the same vanishing condition should also be related to our elliptic Littlewood identity, and indeed we have been able to formulate two conjectures along those lines, Conjectures 5 and 8, which again hold in a number of special cases, and have three different results of [9] as limiting cases.

Macdonald also gave a dual version of the Littlewood identity, in which rather than summing over partitions with even multiplicities, one sums over partitions with even parts. This dual Littlewood identity can, of course, be obtained from the usual Littlewood identity by simply applying Macdonald's involution to conjugate the partitions involved. One can, of course, do the same for the elliptic Littlewood identities, but a new behavior arises. For the μ^2 -type Littlewood identity, there is an analytical symmetry between the parameters p (specifying an elliptic curve) and q (specifying a point on that curve), which is broken by duality. If one attempts to restore this symmetry after dualizing, one finds that, in contrast to the μ^2 -type Littlewood identity, which is a product of two equivalent identities, one p -elliptic, and one q -elliptic, the restoration of symmetry in the dual identity requires that one multiply by a conjectural q -elliptic identity which is *not* equivalent to the original dual identity. Moreover, this partner identity itself has a different broken symmetry, namely the natural action of $SL_2(\mathbb{Z})$ on the family of elliptic curves. One thus finds that each of our identities and conjectures leads to a whole family of conjectures in this way; the Littlewood identity itself gives rise to three conjectural integral transforms, while the other vanishing conjectures correspond to seven different integral transforms. The latter group of conjectures (a single orbit under the various formal symmetries) is particularly interesting, as even without the interpolation functions in the integrands, they would give rise to new transformations of higher-order elliptic

Selberg integrals (specifically, quadratic transformations).

The plan of the paper is as follows. After a discussion of notation at the end of this introduction, we proceed in Section 2 to define our skew interpolation functions, and discuss a number of their properties, especially their connection to ordinary interpolation functions. (We also state a conjectural transformation of higher-order elliptic Selberg integrals related to one of those properties, largely because the same conjecture arose in a different context while working on [8].) Then in Section 3, we discuss the corresponding analogues of the Cauchy identity, along with some necessary preliminaries concerning when skew interpolation functions can be guaranteed to vanish, thus making the relevant sums finite. Section 4 gives the two main forms of the elliptic Littlewood identity, as well as the three associated conjectures at the integral level. Finally, Section 5 discusses a number of conjectures related to the vanishing integrals of [9], with sketches of proofs of various special cases.

The author would like to thank V. Spiridonov and O. Warnaar for helpful comments on an earlier draft, along with R. Askey and M. Rahman for helpful discussions regarding the analogue of Corollary 4.9 for Askey-Wilson polynomials. The author would also like to thank P. Forrester for hosting the author's sabbatical at the University of Melbourne, during which several key portions of the work were done. This work was supported in part by NSF Grant No. DMS-0401387; in addition, much of the work was performed while the author was affiliated with the University of California at Davis.

Notation. We use the notation of [7] and [8]. In particular, bold-face greek letters refer to pairs of partitions; if only one of the partitions is nonzero, we will either give the partition pair explicitly, or rewrite using the notation of [7], explicitly breaking the symmetry between p and q . Thus, for instance, the interpolation functions are denoted by

$$\mathcal{R}_{\lambda}^{*(n)}(z_1, \dots, z_n; a, b; t; p, q), \quad (1.6)$$

which factors as

$$\mathcal{R}_{\lambda, \mu}^{*(n)}(z_1, \dots, z_n; a, b; t; p, q) = R_{\lambda}^{*(n)}(z_1, \dots, z_n; a, b; p, t; q) R_{\mu}^{*(n)}(z_1, \dots, z_n; a, b; q, t; p), \quad (1.7)$$

with the first factor q -elliptic, and the second p -elliptic.

We specifically recall the elliptic Gamma function

$$\Gamma_{p,q}(z) := \prod_{0 \leq i,j} \frac{1 - p^{i+1} q^{j+1} / z}{1 - p^i q^j z}, \quad (1.8)$$

with the convention here (and for Γ^+ , θ , etc.) that multiple arguments express a product:

$$\Gamma_{p,q}(z_1, \dots, z_n) = \prod_{1 \leq i \leq n} \Gamma_{p,q}(z_i). \quad (1.9)$$

This satisfies the functional equations

$$\Gamma_{p,q}(qz) = \theta_p(z) \Gamma_{p,q}(z) \quad (1.10)$$

$$\Gamma_{p,q}(pz) = \theta_q(z) \Gamma_{p,q}(z) \quad (1.11)$$

$$\Gamma_{p,q}(pq/z) = \Gamma_{p,q}(z)^{-1}, \quad (1.12)$$

where

$$\theta_p(z) := \prod_{0 \leq i} (1 - p^i z)(1 - p^{i+1}/z) \quad (1.13)$$

is a theta function, as well as the “quadratic” functional equations

$$\Gamma_{p,q}(z) = \Gamma_{p,q^2}(z, qz) \quad (1.14)$$

$$\Gamma_{p^2,q^2}(z^2) = \Gamma_{p,q}(z, -z), \quad (1.15)$$

which will be useful in the sequel. The special values

$$\Gamma_{p,q^2}(q) = \frac{1}{(q; q^2)} = \frac{(q; q)}{(q^2; q^2)} = (-q; q) \quad (1.16)$$

$$\Gamma_{p,q}(-1) = \frac{(p; p^2)(q; q^2)}{2} \quad (1.17)$$

$$\lim_{x \rightarrow 1} (1-x) \Gamma_{p,q}(x) = \frac{1}{(p; p)(q; q)} \quad (1.18)$$

will also arise. We will also need a third-order elliptic Gamma function

$$\Gamma_{p,q,t}^+(z) := \prod_{0 \leq i,j,k} (1 - p^{i+1} q^{j+1} t^{k+1}/z)(1 - p^i q^j t^k z), \quad (1.19)$$

with functional equations

$$\Gamma_{p,q,t}^+(tz) = \Gamma_{p,q}(z) \Gamma_{p,q,t}^+(z), \quad (1.20)$$

$$\Gamma_{p,q,t}^+(pqt/z) = \Gamma_{p,q,t}^+(z), \quad (1.21)$$

and so forth. (This will only be used to simplify notation; in all of the cases in which it arises, it will appear only via a ratio that resolves via the first functional equation into a product of usual elliptic Gamma functions.)

The elliptic Selberg integral (introduced as the “elliptic Macdonald-Morris conjecture” in [2], and renamed the “Type II” integral in the follow-up [3]) is the integral with density

$$\begin{aligned} \Delta^{(n)}(z_1, \dots, z_n; u_0, \dots, u_5; t; p, q) \\ := \frac{((p; p)(q; q) \Gamma_{p,q}(t/2))^n}{n!} \prod_{1 \leq i < j \leq n} \frac{\Gamma_{p,q}(t z_i^{\pm 1} z_j^{\pm 1})}{\Gamma_{p,q}(z_i^{\pm 1} z_j^{\pm 1})} \prod_{1 \leq i \leq n} \frac{\prod_{0 \leq r < 6} \Gamma_{p,q}(u_r z_i^{\pm 1})}{\Gamma_{p,q}(z_i^{\pm 2})} \frac{dz_i}{2\pi \sqrt{-1} z_i} \end{aligned} \quad (1.22)$$

with associated evaluation ([8], conjectured in [2])

$$\int_{C^n} \Delta^{(n)}(z_1, \dots, z_n; u_0, \dots, u_5; t; p, q) = \prod_{0 \leq i < n} \Gamma_{p,q}(t^{i+1}) \prod_{0 \leq r < s < 6} \Gamma_{p,q}(t^i u_r u_s), \quad (1.23)$$

where the parameters satisfy the *balancing condition*

$$t^{2n-2} \prod_{0 \leq r < 6} u_r = pq, \quad (1.24)$$

and C is a contour such that $C = C^{-1}$, and C contains the rescaled contour tC together with all points of the form $u_r p^i q^j$. (If one allows suitable disjoint unions of contours, this condition can be satisfied unless

$u_r u_s p^i q^j t^k = 1$ for some $0 \leq i, j, k, 0 \leq r, s < 6$.) By convention, the argument $uz_i^{\pm 1}$ to a function indicates a *pair* of arguments $uz_i, u/z_i$, and similarly for $tz_i^{\pm 1} z_j^{\pm 1}$, etc., so in particular the above integrand is hyperoctahedrally symmetric. This determines a natural normalized linear functional

$$\langle f \rangle_{u_0, \dots, u_5; t; p, q}^{(n)} \propto \int_{C^n} f(z_1, \dots, z_n) \Delta^{(n)}(z_1, \dots, z_n; u_0, \dots, u_5; t; p, q), \quad (1.25)$$

where f is a product of p - and q -elliptic functions such that for some nonnegative integers l_r, m_r , the function

$$f(z_1, \dots, z_n) \prod_{\substack{1 \leq i \leq n \\ 0 \leq r \leq 5}} \frac{\Gamma_{p,q}(u_r z_i^{\pm 1})}{\Gamma_{p,q}(p^{-l_r} q^{-m_r} u_r z_i^{\pm 1})} \quad (1.26)$$

is holomorphic, and the contour satisfies the conditions appropriate to

$$\Delta^{(n)}(z_1, \dots, z_n; p^{-l_0} q^{-m_0} u_0, \dots, p^{-l_5} q^{-m_5} u_5; t; p, q); \quad (1.27)$$

the integral is normalized so that

$$\langle 1 \rangle_{u_0, \dots, u_5; t; p, q}^{(n)} = 1. \quad (1.28)$$

Note that if the contour satisfies the conditions for a given choice of l_r, m_r , it satisfies them for all smaller choices, so for a finite linear combination of such functions, one can (generically) choose a contour valid for each term simultaneously, giving linearity. However, the families of functions we consider involve unbounded values of l_0, m_0 , and thus one cannot simply fix a single contour for *every* function in the family. The biorthogonal functions

$$\tilde{\mathcal{R}}_{\lambda}^{(n)}(z_1, \dots, z_n; t_0; t_1, t_2, t_3; u_0, u_1; t; p, q) \quad (1.29)$$

of [8, 7] satisfy biorthogonality with respect to this linear functional; i.e.,

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(z_1, \dots, z_n; t_0; t_1, t_2, t_3; u_0, u_1; t; p, q) \tilde{\mathcal{R}}_{\mu}^{(n)}(z_1, \dots, z_n; t_0; t_1, t_2, t_3; u_1, u_0; t; p, q) \rangle_{t_0, t_1, t_2, t_3, u_0, u_1; t; p, q} \quad (1.30)$$

vanishes unless $\lambda = \mu$.

We will also need higher order versions of the elliptic Selberg integral; we define

$$II_n^{(m)}(u_0, \dots, u_{2m+5}; t; p, q) := \int_{C^n} \Delta^{(n)}(z_1, \dots, z_n; u_0, \dots, u_{2m+5}; t; p, q), \quad (1.31)$$

subject to the balancing condition

$$t^{2n-2} \prod_{0 \leq r < 2m+6} u_r = (pq)^{m+1}, \quad (1.32)$$

in which the density is obtained from the original density ($m = 0$) by replacing

$$\prod_{0 \leq r < 6} \Gamma_{p,q}(u_r z_i^{\pm 1}) \mapsto \prod_{0 \leq r < 2m+6} \Gamma_{p,q}(u_r z_i^{\pm 1}), \quad (1.33)$$

and the contour condition is extended in the obvious way. In particular, if $u_{2m+4} u_{2m+5} = pq$, then the reflection equation for $\Gamma_{p,q}$ causes the two corresponding factors to cancel, reducing m by 1. When $n = 1$, the higher-order elliptic Selberg integral is essentially independent of t , apart from the factor $\Gamma_{p,q}(t)$; we thus define the higher-order elliptic beta integral [13] by

$$I^{(m)}(u_0, \dots, u_{2m+5}; p, q) := \Gamma_{p,q}(t)^{-1} II_1^{(m)}(u_0, \dots, u_{2m+5}; t; p, q); \quad (1.34)$$

note that the constraint that C contains tC is irrelevant in this case.

The factors

$$\Delta_{\lambda}^0(a|b_0, \dots, b_{n-1}; t; p, q) \quad \text{and} \quad \Delta_{\lambda}(a|b_0, \dots, b_{n-1}; t; p, q) \quad (1.35)$$

that appear below are certain multivariate q -symbols (see the introduction of [8]). The first is defined by

$$\Delta_{\lambda}^0(a|b_0, \dots, b_{n-1}; t; p, q) = \prod_{0 \leq r < n} \prod_{1 \leq i \leq \ell(\lambda)} \frac{\theta_p(t^{1-i}(p, q)^{\lambda_i} b_r)}{\theta_p(t^{1-i}(p, q)^{\lambda_i} p q a / b_r)}, \quad (1.36)$$

where $(p, q)^{l, m} := p^l q^m$. The other Δ -symbol is more complicated; the key properties are that the λ -dependent factor of the residue of the elliptic Selberg integrand $\Delta^{(n)}$ at the point $(\dots, (p, q)^{\lambda_i} t^{n-i} u_0, \dots)$ is

$$\Delta_{\lambda}(t^{2n-2} u_0^2 | t^n, t^{n-1} u_0 u_1, \dots, t^{n-1} u_0 u_{2m+5}; t; p, q), \quad (1.37)$$

and that it factors

$$\Delta_{\lambda}(a|b_0, \dots, b_{n-1}; t; p, q) = \Delta_{\lambda}(a; t; p, q) \Delta_{\lambda}^0(a|b_0, \dots, b_{n-1}; t; p, q). \quad (1.38)$$

2 Skew interpolation functions

Consider the following generalized elliptic hypergeometric sum:

$$\begin{aligned} \mathcal{R}_{\lambda/\kappa}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) := & \sum_{\kappa \subset \mu \subset \lambda} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[a/b, ab/pq]; t; p, q} \left\langle \begin{matrix} \mu \\ \kappa \end{matrix} \right\rangle_{[pq/b^2, pq \prod_{0 \leq r < 2n} v_r / ab]; t; p, q} \\ & \Delta_{\mu}^0(pq/b^2 | pq/bv_0, pq/bv_1, \dots, pq/bv_{2n-1}; t; p, q); \end{aligned} \quad (2.1)$$

as the notation suggests, this will turn out to be our desired skew version of the interpolation functions. Note that each term in the rescaled sum

$$\begin{aligned} \hat{\mathcal{R}}_{\lambda/\kappa}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) := & \\ & \frac{\Delta_{\kappa}^0(a/b \prod_{0 \leq r < 2n} v_r | ab/pq \prod_{0 \leq r < 2n} v_r; t; p, q)}{\Delta_{\lambda}^0(a/b | ab/pq; t; p, q)} \mathcal{R}_{\lambda/\kappa}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) \end{aligned} \quad (2.2)$$

is the product of p -abelian and q -abelian factors, so the same applies to this rescaled sum; however, the rescaling introduces unfortunate poles, so we will prefer to use the not-quite-abelian form unless that would introduce complicated factors from quasiperiodicity. This is a generalized elliptic hypergeometric sum in the same sense as the identities of [7]; in particular, it includes the following very-well-poised, balanced, and terminating multivariate elliptic hypergeometric series as a special case:

$$\begin{aligned} \hat{\mathcal{R}}_{(l, m)^n/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) = & \\ \sum_{\mu \subset (l, m)^n} \Delta_{\mu}^0(pq/b^2 | t^n, p^{-l} q^{-m}, p^l q^m a / t^{n-1} b, pq/bv_0, pq/bv_1, \dots, pq/bv_{2n-1}, pq \prod_{0 \leq r < 2n} v_r / ab; t; p, q). \end{aligned} \quad (2.3)$$

We note that the skew interpolation function is invariant under permutations of its arguments, as well as under insertion or deletion of pairs $x, 1/x$. In particular, the arguments are not directly arguments of interpolation functions, but play a more plethystic role. Roughly speaking, this corresponds to the plethystic substitution

$$p_k \mapsto \sum_{0 \leq r < 2n} \frac{v_r^k - v_r^{-k}}{t^{k/2} - t^{-k/2}} \quad (2.4)$$

at the trigonometric level, so that an ordinary argument corresponds to a pair $t^{1/2}x, t^{1/2}/x$ of plethystic arguments. (Compare Theorem 2.5 below.)

The two main identities of [7] both involved sums of this form, and thus one has the following.

Proposition 2.1. [7, Cor. 4.3] *With no arguments, the skew interpolation function is a delta function:*

$$\mathcal{R}_{\lambda/\kappa}([\]; a, b; t; p, q) = \delta_{\lambda\kappa}. \quad (2.5)$$

Proposition 2.2. [7, Thm. 4.1] *With two arguments, the skew interpolation function is an elliptic binomial coefficient:*

$$\mathcal{R}_{\lambda/\kappa}([v_0, v_1]; a, b; t; p, q) = \left\langle \frac{\lambda}{\kappa} \right\rangle_{[a/b, v_0 v_1](a/v_0, a/v_1); t; p, q}. \quad (2.6)$$

Proposition 2.3. [7, Thm. 4.9, Cor. 4.11] *With four arguments, the skew interpolation function has the alternate expressions*

$$\mathcal{R}_{\lambda/\kappa}([v_0, v_1, v_2, v_3]; a, b; t; p, q) = \sum_{\mu} \left\langle \frac{\lambda}{\mu} \right\rangle_{[a/b, v_0 v_1](a/v_0, a/v_1); t; p, q} \left\langle \frac{\mu}{\kappa} \right\rangle_{[a/v_0 v_1 b, v_2 v_3](a/v_0 v_1 v_2, a/v_0 v_1 v_3); t; p, q} \quad (2.7)$$

and

$$\begin{aligned} \mathcal{R}_{\lambda/\kappa}([v_0, v_1, v_2, v_3]; a, b; t; p, q) &= \frac{\Delta_{\lambda}^0(a/b|a/v_0, a/v_1, a/v_2, a/v_3; t; p, q)}{\Delta_{\kappa}^0(a/bV|av_0/V, av_1/V, av_2/V, av_3/V; t; p, q)} \\ &\quad \sum_{\mu} \frac{\left\langle \frac{\lambda}{\mu} \right\rangle_{[a/b, pqV/ab]; t; p, q} \left\langle \frac{\mu}{\kappa} \right\rangle_{[a^2/pqV, ab/pq]; t; p, q}}{\Delta_{\mu}^0(a^2/pqV|a/v_0, a/v_1, a/v_2, a/v_3; t; p, q)}, \end{aligned} \quad (2.8)$$

where $V = v_0 v_1 v_2 v_3$.

In equation (2.7), the binomial coefficients can be expressed in skew interpolation functions, giving

$$\mathcal{R}_{\lambda/\kappa}([v_0, v_1, v_2, v_3]; a, b; t; p, q) = \sum_{\mu} \mathcal{R}_{\lambda/\mu}([v_0, v_1]; a, b; t; p, q) \mathcal{R}_{\mu/\kappa}([v_2, v_3]; a/v_0 v_1, b; t; p, q). \quad (2.9)$$

This generalizes considerably.

Proposition 2.4. *The skew interpolation functions satisfy the identity*

$$\begin{aligned} \mathcal{R}_{\lambda/\kappa}([v_0, \dots, v_{2k-1}, w_0, \dots, w_{2l-1}]; a, b; t; p, q) &= \sum_{\mu} \mathcal{R}_{\lambda/\mu}([v_0, \dots, v_{2k-1}]; a, b; t; p, q) \\ &\quad \mathcal{R}_{\mu/\kappa}([w_0, \dots, w_{2l-1}]; a/v_0 \cdots v_{2k-1}, b; t; p, q). \end{aligned} \quad (2.10)$$

Proof. If we expand the skew interpolation functions on the right via the definition, the inner sum over μ is itself a skew interpolation function with no arguments, and thus the inner sum collapses as required. $\square$

Thus to justify the name “skew interpolation function”, it remains only to show that when $\kappa = 0$, we obtain (a generalization of) the usual interpolation function.

Theorem 2.5. *The interpolation functions have the expression*

$$\mathcal{R}_{\lambda}^{*(n)}(z_1, \dots, z_n; a, b; t; p, q) = \Delta_{\lambda}^0(t^{n-1}a/b|pqa/tb; t; p, q) \mathcal{R}_{\lambda/0}^*([t^{1/2}z_1^{\pm 1}, \dots, t^{1/2}z_n^{\pm 1}]; t^{n-1/2}a, t^{1/2}b; t; p, q). \quad (2.11)$$

Proof. By the connection coefficient identity [7, Cor. 4.14], we can write

$$\mathcal{R}_{\lambda}^{*(n)}(z_1, \dots, z_n; a, b; t; p, q) = \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[t^{n-1}a/b, t^n ab/pq](pqa/tb); t; p, q} \mathcal{R}_{\mu}^{*(n)}(z_1, \dots, z_n; pq/t^n b, b; t; p, q). \quad (2.12)$$

But the new interpolation functions are of “Cauchy” type, so by [7, Prop. 3.9],

$$\mathcal{R}_{\mu}^{*(n)}(z_1, \dots, z_n; pq/t^n b, b; t; p, q) = \Delta_{\mu}^0(pq/tb^2|pqz_1^{\pm 1}/tb, \dots, pqz_n^{\pm 1}/tb; t; p, q) \quad (2.13)$$

as required. $\square$

Remark 1. This can also be proved by induction via the branching rule [7, Thm. 4.16].

Remark 2. Thus ordinary interpolation functions correspond to the case that the parameters multiply pairwise to t ; similarly, the skew interpolation functions of [1] correspond to the special case in which the parameters multiply pairwise to some general, but fixed, r .

Remark 3. The inverse expansion:

$$\Delta_{\lambda}^0(pq/tb^2|pqz_1^{\pm 1}/tb, \dots, pqz_n^{\pm 1}/tb; t; p, q) = \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[pq/tb^2, pq/t^n ab](pqa/tb); t; p, q} \mathcal{R}_{\mu}^{*(n)}(z_1, \dots, z_n; a, b; t; p, q) \quad (2.14)$$

holds even if $\ell(\lambda) > n$ (assuming generic parameters). Indeed, if k is sufficiently large, so that $n + k \geq \ell(\lambda)$, then one may set $z_{n+i} = t^{-i}a$ in

$$\begin{aligned} & \Delta_{\lambda}^0(pq/tb^2|pqz_1^{\pm 1}/tb, \dots, pqz_{n+k}^{\pm 1}/tb; t; p, q) \\ &= \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[pq/tb^2, pq/t^n ab](pqt^{-k}a/tb); t; p, q} \mathcal{R}_{\mu}^{*(n+k)}(z_1, \dots, z_{n+k}; t^{-k}a, b; t; p, q) \end{aligned} \quad (2.15)$$

to obtain the desired result. This will be useful in the sequel, as products of this form satisfy a number of useful identities. For convenience in notation, we will use the product expression (2.13) to extend the Cauchy-type interpolation functions to the case that the indexing partition has more than n parts, as the above considerations eliminate most of the dangers in such an extension.

With this in mind, we refer to the functions $\mathcal{R}_{\lambda/0}^*$ as *lifted interpolation functions*; these seem to be about as close as one can hope to get to an elliptic analogue of the lifted interpolation polynomials of [6, §6]. These functions have a somewhat surprising additional symmetry.

Proposition 2.6. *The lifted interpolation function $\mathcal{R}_{\lambda/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q)$ is invariant under permutations of the $2n+1$ values*

$$v_0, \dots, v_{2n-1}, a / \prod_{0 \leq r < 2n} v_r. \quad (2.16)$$

Proof. Since

$$\left\langle \begin{matrix} \mu \\ 0 \end{matrix} \right\rangle_{[a,b];t;p,q} = \Delta_{\mu}^0(a|b; t; p, q), \quad (2.17)$$

we have

$$\begin{aligned} \mathcal{R}_{\lambda/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) &= \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[a/b, ab/pq];t;p,q} \\ &\quad \Delta_{\mu}^0(pq/b^2 | pq/bv_0, pq/bv_1, \dots, pq/bv_{2n-1}, pq \prod_{0 \leq r < 2n} v_r / ab; t; p, q), \end{aligned} \quad (2.18)$$

which manifestly has the stated symmetry. $\square$

It follows that the connection coefficient formula of [7] extends, and in a particularly nice form.

Corollary 2.7. *One has the identity*

$$\mathcal{R}_{\lambda/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) = \sum_{\kappa} \mathcal{R}_{\lambda/\kappa}^*([a/V, V/a']; a, b; t; p, q) \mathcal{R}_{\kappa/0}^*([v_0, \dots, v_{2n-1}]; a', b; t; p, q), \quad (2.19)$$

where $V = \prod_{0 \leq r < 2n} v_r$.

Proof. Indeed, this reduces to showing

$$\mathcal{R}_{\lambda/0}^*([a/V, V/a', v_0, \dots, v_{2n-1}]; a, b; t; p, q) = \mathcal{R}_{\lambda/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q), \quad (2.20)$$

and this is simply deletion of the pair $V/a', a'/V$ from the left-hand side, after applying Proposition 2.6. $\square$

We also have duality; this breaks the symmetry between p and q , but will be useful in the sequel.

Proposition 2.8. [7, Cor. 4.4]

$$\mathcal{R}_{(0,\lambda)/(0,\kappa)}^*([v_0, \dots, v_{2k-1}]; a, b; t; p, q) = \mathcal{R}_{(0,\lambda')/(0,\kappa')}^*([v_0, \dots, v_{2k-1}]; a, b/qt; 1/q; p, 1/t) \quad (2.21)$$

The other symmetries do respect the p, q symmetry, but lead to unpleasant scale factors since the skew interpolation functions are not quite elliptic, so we use the $\hat{\mathcal{R}}^*$ variant. In particular, this allows one to prove identities by factoring into p -elliptic and q -elliptic factors, then using ellipticity to restore symmetry before multiplying the identities back together.

Proposition 2.9. [7, (4.10)] *If $\ell(\lambda), \ell(\kappa) \leq n$, then*

$$\begin{aligned} \hat{\mathcal{R}}_{(l,m)^n + \lambda / (l,m)^n + \kappa}^*([\dots, v_r, \dots]; a, b; t; p, q) \\ &= \Delta_{(l,m)^n}^0(pq/b^2 | \dots, pq/bv_r, \dots, Qa/t^{n-1}b, p^2q^2t^{n-1}V/Qab; t; p, q) \\ &\quad \times \left(\frac{\Delta_{\lambda}^0(Q^2a/b | pq t^{n-1}Q, Q^2a/t^{n-1}b, ab/pq, p^2q^2Q/b^2; t; p, q)}{\Delta_{\kappa}^0(Q^2a/Vb | pq t^{n-1}Q, Q^2a/t^{n-1}bV, ab/pqV, p^2q^2Q/b^2; t; p, q)} \right) \\ &\quad \times \hat{\mathcal{R}}_{\lambda/\kappa}^*([\dots, v_r, \dots]; Qa, b/Q; t; p, q), \end{aligned} \quad (2.22)$$

where $V = \prod_r v_r$ and $Q = p^l q^m$.

Proposition 2.10. [7, Cor 4.6] If $\lambda_1, \kappa_1 \leq (l, m)$, then

$$\begin{aligned} \hat{\mathcal{R}}_{(l,m)^n \cdot \lambda / (l,m)^n \cdot \kappa}^*([\dots, v_r, \dots]; a, b; t; p, q) \\ = \Delta_{(l,m)^n}^0(pq/b^2 | \dots, pq/bv_r, \dots, Qa/t^{n-1}b, p^2q^2t^{n-1}V/Qab; t; p, q) \\ \times \left(\frac{\Delta_{\lambda}^0(a/t^{2n}b|pq/Qt^{n+1}, Qa/t^{2n-1}b, ab/pq, p^2q^2/t^n b^2; t; p, q)}{\Delta_{\kappa}^0(a/t^{2n}Vb|pq/Qt^{n+1}, Qa/t^{2n-1}bV, ab/pqV, p^2q^2/t^n b^2; t; p, q)} \right) \\ \times \hat{\mathcal{R}}_{\lambda/\kappa}^*([\dots v_r \dots]; t^{-n}a, t^n b; t; p, q). \end{aligned} \quad (2.23)$$

Proposition 2.11. [7, Cor. 4.7] If $\lambda, \kappa \subset (l, m)^n$, then

$$\begin{aligned} \hat{\mathcal{R}}_{(l,m)^n - \kappa / (l,m)^n - \lambda}^*([\dots, v_r, \dots]; a, b; t; p, q) \\ = \Delta_{(l,m)^n}^0(pq/b^2 | \dots, pq/bv_r, \dots, Qa/t^{n-1}b, p^2q^2t^{n-1}V/Qab; t; p, q) \\ \times \left(\frac{\Delta_{\lambda}(t^{2n-2}bV/Q^2a|t^n, 1/Q, t^{n-1}b^2/pqQ, pqV/ab; t; p, q)}{\Delta_{\kappa}(t^{2n-2}b/Q^2a|t^n, 1/Q, t^{n-1}b^2/pqQ, pq/ab; t; p, q)} \right) \\ \times \hat{\mathcal{R}}_{\lambda/\kappa}^*([\dots, v_r, \dots]; pqVt^{n-1}/Qa, pqQ/t^{n-1}b; t; p, q). \end{aligned} \quad (2.24)$$

The above symmetries each follow by applying the corresponding symmetries of elliptic binomial coefficients and Δ symbols to the definition of the skew interpolation functions. There is also an analogue of [7, Cor. 4.8], but this is more subtle. We give this in a fairly general form, for ease of induction and later application.

Theorem 2.12. Suppose the parameters $v_0, \dots, v_{2k-1}$ can be ordered in such a way that $v_{2r}v_{2r+1} = t^{n_r}$ with $n_r \in \mathbb{Z}_{\geq 0}$, $0 \leq r < k$, and let l, m, n, n' be nonnegative integers with $n' = n + \sum_r n_r$ and $\lambda_1, \kappa_1 \leq (l, m)$. Then

$$\begin{aligned} \hat{\mathcal{R}}_{(l,m)^{n'} \cdot \lambda / (l,m)^n \cdot \kappa}^*([v_0, \dots, v_{2k-1}]; a, b; t; p, q) \\ = \frac{\prod_{0 \leq r < k} \Delta_{(l,m)^{n_r}}^0(a/bt^{n'-n_r}|av_{2r}/t^{n'}, av_{2r+1}/t^{n'}; t; p, q)}{\Delta_{(l,m)^{n'-n}}^0(a/bt^n|ab/pq, pqa/t^{n'+n}b; t; p, q)} \\ \times \left(\frac{\Delta_{\kappa}(a/t^{n'+n}b|p^{-l}q^{-m}, p^lq^ma/t^{n'-1}b, ab/pqt^{n'}, pqt^{n'-n}/ab; t; p, q)}{\Delta_{\lambda}(a/t^{2n'}b|p^{-l}q^{-m}, p^lq^ma/t^{n'-1}b, ab/pqt^{n'}, pq/ab; t; p, q)} \right) \\ \times \hat{\mathcal{R}}_{\kappa/\lambda}^*([v_0, \dots, v_{2k-1}]; pq/t^n b, pqt^{n'}/a; t; p, q) \end{aligned} \quad (2.25)$$

Proof. When $k = 1$, this follows immediately from Corollaries 4.6 and 4.8 of [7]. (Corollary 4.8 corresponds to the case $n = 0$, $k = 1$, and Corollary 4.6 allows one to extend this to $n > 0$.) We then proceed by induction on k . One first notes that

$$\begin{aligned} \hat{\mathcal{R}}_{(l,m)^{n'} \cdot \lambda / (l,m)^n \cdot \kappa}^*([v_0, \dots, v_{2k-1}]; a, b; t; p, q) \\ = \sum_{\mu} \hat{\mathcal{R}}_{(l,m)^{n'} \cdot \lambda / \mu}^*([v_0, v_1]; a, b; t; p, q) \hat{\mathcal{R}}_{\mu / (l,m)^n \cdot \kappa}^*([v_2, \dots, v_{2k-1}]; a/t^{n_0}, b; t; p, q). \end{aligned} \quad (2.26)$$

The key observation is that the first factor is

$$\left\langle \begin{matrix} (l, m)^{n'} \cdot \lambda \\ \mu \end{matrix} \right\rangle_{[a/b, t^{n_0}](a/v_0, a/v_1, p^2q^2/b^2); t; p, q}, \quad (2.27)$$

which vanishes unless

$$\boldsymbol{\mu}_i \leq ((l, m)^{n'} \cdot \boldsymbol{\lambda})_i \leq \boldsymbol{\mu}_{i-n_0}. \quad (2.28)$$

In particular, for $1 \leq i \leq n'$, $\boldsymbol{\mu}_i \leq (l, m)$, while for $1 \leq i \leq n' - n_0$, $\boldsymbol{\mu}_i \geq (l, m)$. We can thus rewrite the sum as

$$\sum_{\boldsymbol{\mu}} \hat{\mathcal{R}}_{(l, m)^{n'} \cdot \boldsymbol{\lambda} / (l, m)^{n'} - n_0 \cdot \boldsymbol{\mu}}^*([v_0, v_1]; a, b; t; p, q) \hat{\mathcal{R}}_{(l, m)^{n'} - n_0 \cdot \boldsymbol{\mu} / (l, m)^n \cdot \boldsymbol{\kappa}}^*([v_2, \dots, v_{2k-1}]; a/t^{n_0}, b; t; p, q). \quad (2.29)$$

The result follows by applying the symmetry to each factor and simplifying. $\square$

Dually, one has the following identity.

Corollary 2.13. *Suppose the parameters $v_0, \dots, v_{2k-1}$ can be ordered in such a way that $v_{2r}v_{2r+1} = Q_r^{-1}$, $Q_r := p^{l_r}q^{m_r}$, with $l_r, m_r \in \mathbb{Z}_{\geq 0}$ for $0 \leq r < k$, and let l, l', m, m', n , be nonnegative integers with $l' = l + \sum_r l_r$, $m' = m + \sum_r m_r$, $\ell(\boldsymbol{\lambda}), \ell(\boldsymbol{\kappa}) \leq n$. Then*

$$\begin{aligned} \hat{\mathcal{R}}_{(l', m')^n + \boldsymbol{\lambda} / (l, m)^n + \boldsymbol{\kappa}}^*([v_0, \dots, v_{2k-1}]; a, b; t; p, q) \\ = \frac{\prod_{0 \leq r < k} \Delta_{(l_r, m_r)^n}^0(Q' a / Q_r b | Q' a / Q_r v_{2r}, Q' a / Q_r v_{2r+1}; t; p, q)}{\Delta_{(l' - l, m' - m)^n}^0(Q a / b | a b / p q, p q Q Q' a / b; t; p, q)} \\ \times \left(\frac{\Delta_{\boldsymbol{\kappa}}(a Q Q' / b | t^n, a t Q' / t^n b, a b Q' / p q, p q Q / Q' a b; t; p, q)}{\Delta_{\boldsymbol{\lambda}}(a Q'^2 / b | t^n, a t Q' / t^n b, a b Q' / p q, p q / a b; t; p, q)} \right) \\ \times \hat{\mathcal{R}}_{\boldsymbol{\kappa} / \boldsymbol{\lambda}}^*([v_0, \dots, v_{2k-1}]; p q Q / b, p q / Q' a; t; p, q), \end{aligned} \quad (2.30)$$

where $Q = p^l q^m$, $Q' = p^{l'} q^{m'}$.

If $\boldsymbol{\lambda} = 0$ in the first identity, one can apply complementation to obtain a relation between $\hat{\mathcal{R}}_{\boldsymbol{\kappa}/0}^*$ and $\hat{\mathcal{R}}_{(l, m)^n - \boldsymbol{\kappa}/0}^*$; the constraint on the arguments causes both of these to be ordinary interpolation functions in n variables, and this is just the usual complementation symmetry of such functions. (In contrast, in the corresponding special case of the corollary, the lifted interpolation functions are not simply ordinary interpolation functions.) Particularly interesting is the case that both $\boldsymbol{\kappa}$ and its complement are rectangles, since then the identity is a transformation of more classically hypergeometric sums (under the hypotheses of Theorem 2.12):

$$\begin{aligned} \hat{\mathcal{R}}_{(l, m)^n / 0}^*([v_0, \dots, v_{2k-1}]; a, b; t; p, q) \\ \propto \prod_{0 \leq i < 2k} \frac{\Gamma_{p, q, t}^+(b/v_i, (Q(pqt/b))/v_i, (t^{-n-n'}at)/v_i, (pq/Qt^{-n-n'}a)/v_i)}{\Gamma_{p, q, t}^+(bv_i, Q(pqt/b)v_i, t^{-n-n'}atv_i, (pqt/Qt^{-n-n'}at)v_i)} \\ \hat{\mathcal{R}}_{(l, m)^{n'} / 0}^*([v_0, \dots, v_{2k-1}]; t^{n+n'-1}b/Q, Qa/t^{n+n'-1}; t; p, q), \end{aligned} \quad (2.31)$$

where the constant of proportionality can be determined from the case $k = 1$, when both lifted interpolation functions have explicit evaluations. This is a sort of dual Karlsson-Minton sum; in particular, the dual of this sum (coming from the Corollary) is a multivariate analogue of [11, Cor. 4.5].

As usual with such sums, there appears to be an integral analogue of (2.31).

Conjecture 1. For integers $m, n, n_0, \dots, n_{k-1} \geq 0$, and parameters $t_0, t_1, t_2, t_3, v_0, \dots, v_{2k-1}$ satisfying

$$t_0 t_1 t_2 t_3 = t^{2+m-n} \quad (2.32)$$

$$v_{2i} v_{2i+1} = pq/t^{n_i} \quad (2.33)$$

$$\sum_{0 \leq i < k} n_i = m + n, \quad (2.34)$$

one has

$$\begin{aligned} \Pi_n^{(k-1)}(t_0, t_1, t_2, t_3, v_0, \dots, v_{2k-1}; t; p, q) &= \prod_{m < i \leq n} \prod_{0 \leq r < s < 4} \Gamma_{p,q}(t^{n-i} t_r t_s) \prod_{0 \leq i < 2k} \prod_{0 \leq r < 4} \frac{\Gamma_{p,q,t}^+(p q t_r / v_i)}{\Gamma_{p,q,t}^+(t_r v_i)} \\ \Pi_m^{(k-1)}(t/t_0, t/t_1, t/t_2, t/t_3, v_0, \dots, v_{2k-1}; t; p, q). \end{aligned} \quad (2.35)$$

This holds when $k = 1$ (both sides can be explicitly evaluated), as well as when $k = 2$, as a special case of the E_8 symmetry of [8] (a rare case in which a transformation outside the usual double cosets can be applied, via a sequence of two dimension-changing transformations). The case $t \mapsto pq/t$, $|m - n| \leq 1$ appears naturally if one attempts to give a direct proof of the commutation relations for the integral operators of [8]. (Note that the case $m = n$ implies the general case, as one may take the limit $v_0 \rightarrow t^{-n_0} t_3$ to reduce the dimension on the right-hand side.)

3 Elliptic Cauchy identities

Thus, in particular, the skew interpolation functions behave very much as analogues of skew Macdonald polynomials—not entirely surprising, given that skew Macdonald polynomials are limits of skew interpolation functions, as follows from Theorem 8.5 of [7]. More precisely, one has

$$\begin{aligned} \lim_{p \rightarrow 0} p^{|\lambda|/4 - |\mu|/4} \mathcal{R}_{(0,\lambda)/(0,\mu)}^*([p^{1/4}/v_0, \dots, p^{1/4}/v_{n-1}, p^{-1/4}w_0, \dots, p^{-1/4}w_{n-1}]; a, p^{1/2}b; t; p, q) \\ = \frac{(-a)^{|\lambda|} q^{n(\lambda')} t^{-2n(\lambda)} C_{\lambda}^-(t; q, t)}{(-aV/W)^{|\mu|} q^{n(\mu')} t^{-2n(\mu)} C_{\mu}^-(t; q, t)} P_{\lambda/\mu}([\frac{v_0^k + \dots + v_{n-1}^k - w_0^k - \dots - w_{n-1}^k}{1 - t^k}]; q, t), \end{aligned} \quad (3.1)$$

by a straightforward induction from the case $l = 1$.

However, if we attempt to give a direct analogue of the Cauchy identity, we encounter the difficulty that sums of infinitely many elliptic terms rarely converge. It will thus be important to understand under what circumstances a skew interpolation function is forced to vanish.

Lemma 3.1. If λ, κ are partition pairs, l, m, n nonnegative integers, and a, b , and $v_0 \in \mathbb{C}^*$ are generic, then

$$\mathcal{R}_{\lambda/\kappa}^*([v_0, t^n/p^l q^m v_0]; a, b; t; p, q) \quad (3.2)$$

vanishes unless

$$\kappa_i \leq \lambda_i \leq \kappa_{i-n} + (l, m) \quad (3.3)$$

for all i , with the convention $\kappa_0 = \kappa_{-1} = \dots = \infty$.

Proof. Observe that we can write

$$\begin{aligned} \mathcal{R}_{\lambda/\kappa}^*([v_0, t^n/p^l q^m v_0]; a, b; t; p, q) &= \sum_{\mu} \mathcal{R}_{\lambda/\mu}^*([v_0, t^n/v_0]; a, b; t; p, q) \\ &\quad \mathcal{R}_{\mu/\kappa}^*([v_0/t^n, t^n/p^l q^m v_0]; a/t^n, b; t; p, q), \end{aligned} \quad (3.4)$$

with

$$\mathcal{R}_{\lambda/\mu}^*([v_0, t^n/v_0]; a, b; t; p, q) = \frac{\Delta_{\lambda}^0(a/b|a/v_0, av_0/t^n; t; p, q)}{\Delta_{\mu}^0(a/bt^n|a/v_0, av_0/t^n; t; p, q)} \left\langle \frac{\lambda}{\mu} \right\rangle_{[ab, t^n]; t; p, q}, \quad (3.5)$$

and

$$\mathcal{R}_{\mu/\kappa}^*([v_0/t^n, t^n/p^l q^m v_0]; a/t^n, b; t; p, q) = \frac{\Delta_{\mu}^0(a/bt^n|a/v_0, av_0 p^l q^m/t^{2n}; t; p, q)}{\Delta_{\kappa}^0(ap^l q^m/bt^n|a/v_0, av_0 p^l q^m/t^{2n}; t; p, q)} \left\langle \frac{\mu}{\kappa} \right\rangle_{[a/bt^n, p^{-l} q^{-m}]; t; p, q}. \quad (3.6)$$

The binomial coefficients vanish unless [7, Cor. 4.5]

$$\mu_i \leq \lambda_i \leq \mu_{i-n} \quad (3.7)$$

and [7, Cor. 4.2]

$$\kappa_i \leq \mu_i \leq \kappa_i + (l, m), \quad (3.8)$$

and (by genericity), this vanishing cannot be cancelled by a pole of the remaining factors. $\square$

The other significant source of vanishing is the following.

Lemma 3.2. *If λ, κ are partition pairs, l, m, n are nonnegative integers, a, b , and $v_0 \in \mathbb{C}^*$ are generic, and $\kappa_{n+1} \leq (l, m)$, then*

$$\mathcal{R}_{\lambda/\kappa}^*([v_0, a/p^{-l} q^{-m} t^n]; a, b; t; p, q) \quad (3.9)$$

vanishes unless $\lambda_{n+1} \leq (l, m)$.

Proof. We have

$$\mathcal{R}_{\lambda/\kappa}^*([v_0, a/p^{-l} q^{-m} t^n]; a, b; t; p, q) = \frac{\Delta_{\lambda}^0(a/b|p^{-l} q^{-m} t^n; t; p, q)}{\Delta_{\kappa}^0(p^{-l} q^{-m} t^n/bv_0|p^{-l} q^{-m} t^n; t; p, q)} \left\langle \frac{\lambda}{\kappa} \right\rangle_{[a/b, av_0/p^{-l} q^{-m} t^n](v_0); t; p, q}. \quad (3.10)$$

The binomial coefficient factor is generic, so cannot contribute any poles, as are the factors coming from denominators of Δ^0 . We are thus left with considering the ratio

$$\frac{\mathcal{C}_{\lambda}^0(p^{-l} q^{-m} t^n; t; p, q)}{\mathcal{C}_{\kappa}^0(p^{-l} q^{-m} t^n; t; p, q)} \quad (3.11)$$

If $\kappa_{n+1} \leq (l, m)$, then the denominator is nonzero; the numerator vanishes unless $\lambda_{n+1} \leq (l, m)$. $\square$

Theorem 3.3. *Suppose the sequence $v_0, v_1, \dots, v_{2k-1}$ can be ordered in such a way that $v_{2i} v_{2i+1} = t^{n_i} p^{-l_i} q^{-m_i}$ with $l_i, m_i, n_i \geq 0$, for $0 \leq i < k$, and are otherwise generic. Then for any partition pair κ ,*

$$\mathcal{R}_{\lambda/\kappa}^*([v_0, v_1, \dots, v_{2k-1}]; a, b; t; p, q) = 0 \quad (3.12)$$

unless

$$\kappa_i \leq \lambda_i \leq \kappa_{i-N} + (L, M), \quad (3.13)$$

where $L = \sum_i l_i$, $M = \sum_i m_i$, $N = \sum_i n_i$.

Proof. If $k = 1$ or $k = 0$, this follows from Lemma 3.1. In general, we have

$$\mathcal{R}_{\lambda/\kappa}([v_0, v_1, \dots, v_{2k-1}]; a, b; t; p, q) = \sum_{\mu} \mathcal{R}_{\lambda/\mu}([v_0, v_1]; a, b; t; p, q) \mathcal{R}_{\mu/\kappa}([v_2, \dots, v_{2k-1}]; a/v_0 v_1, b; t; p, q); \quad (3.14)$$

the term associated to μ vanishes unless

$$\mu_i \leq \lambda_i \leq \mu_{i-n_0} + (l_0, m_0) \quad (3.15)$$

and (by induction)

$$\kappa_i \leq \mu_i \leq \kappa_{i-(N-n_0)} + (L - l_0, M - m_0). \quad (3.16)$$

The claim follows. $\square$

Similarly, one has the following.

Theorem 3.4. *Let $l_0, \dots, l_{k-1}$, $m_0, \dots, m_{k-1}$, $n_0, \dots, n_{k-1}$ be sequences of nonnegative integers, and suppose the otherwise generic sequence $v_0, \dots, v_{2k-1}$ can be ordered in such a way that $v_{2i} v_{2i+1} = t^{n_i} p^{-l_i} q^{-m_i}$ for $1 \leq i < k$, while*

$$a / \prod_{1 \leq i < 2k} v_i = t^{n_0} p^{-l_0} q^{-m_0}. \quad (3.17)$$

If $\kappa_{n_0+1} \leq (l_0, m_0)$, then

$$\mathcal{R}_{\lambda/\kappa}([v_0, v_1, \dots, v_{2k-1}]; a, b; t; p, q) = 0 \quad (3.18)$$

unless $\lambda_{N+1} \leq (L, M)$.

When $\kappa = 0$, the two vanishing conditions coincide, and both simply state that $\lambda_{N+1} \leq (L, M)$. This corresponds to the extra symmetry explained in Proposition 2.6 above. One also obtains an additional (albeit more delicate) source of vanishing in the $\kappa = 0$ case.

Theorem 3.5. *Let $l \leq L$; $m \leq M$; $n \leq N$ be nonnegative integers. Then the lifted interpolation function*

$$\mathcal{R}_{\lambda/0}([p^L q^M a/t^N, v_1, \dots, v_{2k-1}]; a, pqt^n/p^l q^m a; t; p, q) \quad (3.19)$$

vanishes unless $\lambda_{N+1} \leq (L, M)$.

Proof. We have the expansion

$$\begin{aligned} & \mathcal{R}_{\lambda/0}([p^L q^M a/t^N, v_1, \dots, v_{2n-1}]; a, pqt^n/p^l q^m a; t; p, q) \\ &= \sum_{\mu} \left\langle \frac{\lambda}{\mu} \right\rangle_{[a/b, t^n/p^l q^m]; t; p, q} \Delta_{\mu}^0(pq/b^2 | t^{N-n}/p^{L-l} q^{M-m}, pq/bv_1, \dots, pq/bv_{2n-1}, pq \prod_{0 \leq r < 2n} v_r/ab; t; p, q), \end{aligned} \quad (3.20)$$

where $b = pqt^n/p^l q^m a$. The second factor vanishes unless

$$\mu_{N-n+1} \leq (L-l, M-m), \quad (3.21)$$

while the first factor vanishes unless

$$\mu_i \leq \lambda_i \leq \mu_{i-n} + (l, m). \quad (3.22)$$

□

Since infinite sums of elliptic functions tend not to converge, we need to insist in the elliptic Cauchy identity that the sum terminate; i.e., involve only finitely many terms. To avoid potential obstructions to analytic continuation arguments, we insist that the termination occurs either because the partition pair being summed over occurs as the lower partition in a skew interpolation function (or elliptic binomial coefficient), or because using either Δ^0 factors of the summand or one of the first two vanishing theorems, one can bound both the first part and the length of the partition pair. In the latter case, we will refer to the source of the bound on the first part as a horizontal termination condition; similarly a vertical termination condition is one that allows us to bound the length.

With this in mind, we can now state our first version of the Cauchy identity for skew interpolation functions. Note that the termination conditions allow the right-hand side to be simplified to an expression in p -theta and q -theta functions; this would not hold if the sum were finite by virtue of the third vanishing condition alone.

Theorem 3.6. *One has the identity*

$$\sum_{\mu} \Delta_{\mu}(a/b|; t; p, q) \mathcal{R}_{\mu/0}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) \\ \mathcal{R}_{\mu/0}^*([w_0, \dots, w_{2m-1}]; \sqrt{pqt}/b, \sqrt{pqt}/a; t; p, q) = \prod_{\substack{0 \leq i < 2n+2 \\ 0 \leq j < 2m+2}} \frac{\Gamma_{p,q,t}^+((pqt)^{1/2} v_i / w_j)}{\Gamma_{p,q,t}^+((pqt)^{1/2} v_i w_j)}, \quad (3.23)$$

where

$$v_{2n} = a / \prod_{0 \leq r < 2n} v_r, \quad v_{2n+1} = 1/a, \quad w_{2n} = (pqt)^{1/2}/b \prod_{0 \leq r < 2m} w_r, \quad w_{2n+1} = b/(pqt)^{1/2}, \quad (3.24)$$

and the parameters are such that the sum terminates, but otherwise generic.

Proof. Suppose first that the vertical termination of the sum is due to the v parameters, while the horizontal termination is due to the w parameters. We may thus assume (adding or removing pairs $x, 1/x$ as necessary) that $v_{2i}v_{2i+1} = t$, $0 \leq i < n$, while $w_{2i}w_{2i+1} = 1/p$ or $1/q$ for each $0 \leq i < m$. In that case, we may factor the sum into the product of a q -elliptic sum and a p -elliptic sum. Applying duality to the w factor allows us to express both factors as interpolation functions, and the claim becomes the Cauchy identity of [7, Thm. 4.18].

The other possibility (up to obvious symmetries) is that one set of parameters (say the w parameters) provides both termination conditions. If the v parameters *also* provide vertical termination, then the result follows; in general, the set of v parameters for which $v_{2i}v_{2i+1} \in t^{\mathbb{N}}$, $0 \leq i < n$, is Zariski dense on both elliptic curves, so we may analytically continue. □

There is also a skew version of the above identity.

Theorem 3.7. *One has the identity*

$$\begin{aligned} & \sum_{\mu} \frac{\Delta_{\mu}(a/b|; t; p, q)}{\Delta_{\lambda}(a/bV|; t; p, q)} \mathcal{R}_{\mu/\lambda}^*([v_0, \dots, v_{2n-1}]; a, b; t; p, q) \mathcal{R}_{\mu/\kappa}^*([w_0, \dots, w_{2m-1}]; \sqrt{pqt}/b, \sqrt{pqt}/a; t; p, q) \\ & \propto \sum_{\mu} \mathcal{R}_{\lambda/\mu}^*([w_0, \dots, w_{2m-1}]; \sqrt{pqt}/b, \sqrt{pqt}V/a; t; p, q) \frac{\Delta_{\kappa}(a/bW|; t; p, q)}{\Delta_{\mu}(a/bVW|; t; p, q)} \mathcal{R}_{\kappa/\mu}^*([v_0, \dots, v_{2n-1}]; a, bW; t; p, q), \end{aligned} \quad (3.25)$$

where $V = \prod_r v_r$, $W = \prod_r w_r$, for generic parameters such that the left-hand side terminates. The constant of proportionality is independent of λ and κ , and is thus equal to the value of the sum when $\lambda = \kappa = 0$.

Proof. First consider the case $\kappa = 0$, so that the right-hand side collapses to $\Delta_{\lambda}(a/bV|; t; p, q)$ (times the value of the sum for $\lambda = \kappa = 0$), and suppose furthermore that $v_{2i-1}v_{2i} = t$, and $w_{2i-1}w_{2i} \in p^{-\mathbb{N}}q^{-\mathbb{N}}$ for each i . If we multiply both sides by

$$\mathcal{R}_{\lambda/0}^*([t^{1/2}u_1^{\pm 1}, \dots, t^{1/2}u_{\ell(\lambda)}^{\pm 1}]; a/V, b; t; p, q) \quad (3.26)$$

and sum over λ , the right-hand side becomes an instance of the previous theorem, while the left-hand side simplifies directly to an instance of the previous theorem. In particular, after so multiplying and summing, the two sides agree. But the test functions we have multiplied by are linearly independent, and thus both sides agree before summing.

The arbitrary terminating case with $\kappa = 0$ then follows by analytic continuation. Similarly, the case $\kappa \neq 0$ follows from the case $\kappa = 0$, and the general claim follows by analytic continuation. $\square$

Another approach to proving the above identity is by induction on n and m ; it suffices to consider the case $n = m = 1$, or in other words the following special case.

Corollary 3.8. *One has the identity*

$$\begin{aligned} & \sum_{\mu} \frac{\Delta_{\mu}(a|v_0, v_1, v_2, v_3; t; p, q)}{\Delta_{\lambda}(a/b_0|v_0, v_1, v_2, v_3; t; p, q)} \left\langle \frac{\mu}{\lambda} \right\rangle_{[a, b_0]; t; p, q} \left\langle \frac{\mu}{\kappa} \right\rangle_{[a, b_1]; t; p, q} \\ & \propto \sum_{\mu} \left\langle \frac{\lambda}{\mu} \right\rangle_{[a/b_0, b_1]; t; p, q} \frac{\Delta_{\kappa}(a/b_1|v_0, v_1, v_2, v_3; t; p, q)}{\Delta_{\mu}(a/b_0b_1|v_0, v_1, v_2, v_3; t; p, q)} \left\langle \frac{\kappa}{\mu} \right\rangle_{[a/b_1, b_0]; t; p, q}, \end{aligned} \quad (3.27)$$

assuming the termination conditions

$$t^{\mathbb{N}} \cap \{b_0, b_1, v_0, v_1, v_2, v_3\} \neq \emptyset \quad (3.28)$$

$$p^{-\mathbb{N}}q^{-\mathbb{N}} \cap \{b_0, b_1, v_0, v_1, v_2, v_3\} \neq \emptyset, \quad (3.29)$$

(with corresponding conditions on λ, κ if the v_r are used for termination), and the balancing condition $b_0b_1v_0v_1v_2v_3 = pqta^2$. The constant of proportionality is given by

$$\sum_{\mu} \Delta_{\mu}(a|b_0, b_1, v_0, v_1, v_2, v_3; t; p, q). \quad (3.30)$$

This, in turn, gives an expression for the (discrete) inner product of two interpolation functions with respect to the density of [7, Thm. 5.8]; in particular, it is a limiting case of an integral identity [8, Thm. 9.4]. It can also be obtained (following ideas of [10]) by computing connection coefficients for interpolation theta functions in two different ways (compare [7, Thm. 4.15]).

4 Elliptic Littlewood identities

Since the classical Littlewood identity only involves a single Schur function, the termination conditions in any direct elliptic analogue must be borne by a single skew interpolation function. It turns out, however, that the conditions can be weakened slightly; it is permissible for the skew interpolation function to allow unbounded upper partitions, so long as none of those satisfy the even multiplicity condition. The point is that since the first and second parts of μ^2 agree, we need only have a bound on the *second* part of μ^2 to obtain a terminating sum. Thus in the horizontal termination condition, we may allow one of the pairs to multiply to $tp^{-l_i}q^{-m_i}$ instead of $p^{-l_i}q^{-m_i}$.

In particular, if $v_0v_1 = t$, then this simultaneously gives both horizontal and vertical termination conditions. Indeed, we find that if $\mathcal{R}_{\mu^2/\lambda}^*([v_0, t/v_0]; a, b; t; p, q) \neq 0$, then

$$\lambda_{2i-1} \leq (\mu^2)_{2i-1} \leq \lambda_{2i-2} \quad (4.1)$$

$$\lambda_{2i} \leq (\mu^2)_{2i} \leq \lambda_{2i-1}, \quad (4.2)$$

and thus, since $(\mu^2)_{2i-1} = (\mu^2)_{2i} = \mu_i$,

$$\lambda_{2i-1} \leq \mu_i \leq \lambda_{2i-1}, \quad (4.3)$$

so that μ is uniquely determined. With this in mind, define new operations on partition pairs

$$\lambda^+ : \lambda_i^+ = \lambda_{2i-1} \quad (4.4)$$

$$\lambda^- : \lambda_i^- = \lambda_{2i-2}. \quad (4.5)$$

Lemma 4.1. *For any partition pair λ ,*

$$\begin{aligned} \sum_{\mu} \left\langle \begin{matrix} \mu^2 \\ \lambda \end{matrix} \right\rangle_{[a,t];t;p,q} \Delta_{\mu}(a/t|v_0, \dots, v_{2n-1}; t^2; p, q) \\ = \Delta_{\lambda}(a/t|v_0, \dots, v_{2n-1}; t; p, q) \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu^2 \end{matrix} \right\rangle_{[a/t,t];t;p,q} \frac{\Delta_{\mu}(a/t^3|v_0, \dots, v_{2n-1}; t^2; p, q)}{\Delta_{\mu^2}(a/t^2|v_0, \dots, v_{2n-1}; t; p, q)} \end{aligned} \quad (4.6)$$

Proof. On the left-hand side, only the term with $\mu = \lambda^+$ contributes, while on the right-hand side, only the term with $\mu = \lambda^-$ contributes. Now, it follows easily from the definition of Δ^0 that

$$\Delta_{\lambda}^0(a/t|v_0; t; p, q) = \Delta_{\lambda^+}^0(a/t|v_0; t^2; p, q) \Delta_{\lambda^-}^0(a/t^3|v_0/t; t^2; p, q), \quad (4.7)$$

$$\Delta_{\mu^2}^0(a/t|v_0; t; p, q) = \Delta_{\mu}^0(a/t^2|v_0, v_0/t; t^2; p, q), \quad (4.8)$$

and thus the dependence on x and v_r disappears. It thus suffices to show that

$$\left\langle \begin{matrix} \lambda^{+2} \\ \lambda \end{matrix} \right\rangle_{[a,t];t;p,q} = \left\langle \begin{matrix} \lambda \\ \lambda^{-2} \end{matrix} \right\rangle_{[a/t,t];t;p,q} \frac{\Delta_{\lambda}(a/t; t; p, q) \Delta_{\lambda^-}(a/t^3; t^2; p, q)}{\Delta_{\lambda^{-2}}(a/t^2; t; p, q) \Delta_{\lambda^+}(a/t; t^2; p, q)}. \quad (4.9)$$

This can be proved by induction in $\ell(\lambda)$ via the observations

$$((l, m) \cdot \lambda)^+ = (l, m) \cdot \lambda^- \quad (4.10)$$

$$((l, m) \cdot \lambda)^- = \lambda^+ \quad (4.11)$$

and the relation [7, Cor. 4.8]

$$\left\langle \begin{matrix} (l, m) \cdot \lambda \\ \mu \end{matrix} \right\rangle_{[a, t]; t; p, q} = \Delta_{(l, m)}^0(a|t; t; p, q) \frac{\Delta_{\mu}(a/t|p^{-l}q^{-m}, p^l q^m a; t; p, q)}{\Delta_{\lambda}(a/t^2|p^{-l}q^{-m}, p^l q^m a; t; p, q)} \left\langle \lambda \right\rangle_{[a/t, t]; t; p, q}. \quad (4.12)$$

Note that we may freely check the p -theta and q -theta portions of the relation separately, and rescale so that both are elliptic. $\square$

The first version of the elliptic Littlewood identity is the following.

Theorem 4.2. *We have*

$$\begin{aligned} & \sum_{\mu} \mathcal{R}_{\mu^2/0}^*([v_0, \dots, v_{2n-1}]; ta, (pqt)^{1/2}/a; t; p, q) \Delta_{\mu}(a^2/(pqt)^{1/2}; t^2; p, q) \\ &= \frac{\Gamma_{p, q, t}^+((pqt)^{1/2})^n \Gamma_{p, q, t}^+((pqt)^{1/2}t) \prod_{0 \leq i < j < 2n+2} \Gamma_{p, q, t}^+((pqt)^{1/2}v_i/v_j)}{\prod_{0 \leq i < 2n+2} \Gamma_{p, q, t^2}^+((pqt)^{1/2}v_i^2) \prod_{0 \leq i < j < 2n+2} \Gamma_{p, q, t}^+((pqt)^{1/2}v_i v_j)}, \end{aligned} \quad (4.13)$$

where $v_{2n} = ta / \prod_{0 \leq i < 2n} v_i$, $v_{2n+1} = 1/a$, and the sum terminates.

Proof. Using the S_{2n+1} symmetry of the lifted interpolation functions, we may assume (inserting $x, 1/x$ pairs as necessary) that the parameters pairwise multiply to t , and are ordered in such a way that $v_{2m}, \dots, v_{2n-1}$ gives both horizontal and vertical termination conditions for $0 \leq m < n$. The proof then follows by a straightforward induction in n :

$$\begin{aligned} & \sum_{\mu} \mathcal{R}_{\mu^2/0}^*([v_0, \dots, v_{2n-1}]; ta, b; t; p, q) \Delta_{\mu}(a/b; t^2; p, q) \\ &= \sum_{\lambda} \mathcal{R}_{\lambda/0}^*([v_2, \dots, v_{2n-1}]; a, b; t; p, q) \\ & \quad \sum_{\mu} \mathcal{R}_{\mu^2/\lambda}^*([v_0, t/v_0]; ta, b; t; p, q) \Delta_{\mu}(a/b; t^2; p, q) \end{aligned} \quad (4.14)$$

$$\begin{aligned} &= \sum_{\mu} \frac{\Delta_{\mu}(a/bt^2; t^2; p, q)}{\Delta_{\mu^2}(a/bt; t; p, q)} \\ & \quad \sum_{\lambda} \Delta_{\lambda}(a/b; t; p, q) \mathcal{R}_{\lambda/0}^*([v_2, \dots, v_{2n-1}]; a, b; t; p, q) \mathcal{R}_{\lambda/\mu^2}^*([v_0, t/v_0]; a, b; t; p, q) \end{aligned} \quad (4.15)$$

$$\propto \sum_{\mu} \Delta_{\mu}(a/bt^2; t^2; p, q) \mathcal{R}_{\mu^2/0}^*([v_2, \dots, v_{2n-1}]; a, bt; t; p, q). \quad (4.16)$$

Note that the last step only works if $ab = (pqt)^{1/2}$. $\square$

Remark. Note that the termination condition prevents one from obtaining a Macdonald polynomial identity as a simple limit, except in the case $n = 1$. However, if one ignores the issue of termination, and takes a limit

above, one obtains

$$\sum_{\mu} \frac{C_{\mu}^{-}(t; q, t^2)}{C_{\mu}^{-}(q; q, t^2)} P_{\mu^2} \left(\left[\frac{v_0^k + \dots + v_{n-1}^k - w_0^k - \dots - w_{n-1}^k}{1 - t^k} \right]; q, t \right) \\ = \frac{\prod_{0 \leq i, j < n} (v_i w_j; q, t)}{\prod_{0 \leq i < n} (w_i^2, t v_i^2; q, t^2) \prod_{0 \leq i < j < n} (v_i v_j, w_i w_j; q, t)}, \quad (4.17)$$

agreeing with Macdonald's q, t -Littlewood identity. This agreement results from the fact that the $n = 1$ case and the Cauchy identity together suffice to make the above induction work in the absence of termination. One can presumably also obtain this limit by restricting p to a geometric progression $p = q^{2^m}$, $m \rightarrow \infty$, thus allowing the sum to terminate for all sufficiently large m .

If the lifted interpolation function is terminating in the usual sense (i.e., without taking advantage of the one extra factor of t), then it in fact corresponds to an ordinary interpolation function evaluated at a partition. This gives rise to the following identity.

Corollary 4.3. *For every partition pair λ , one has the following identity of meromorphic functions*

$$\sum_{\mu} \frac{\Delta_{\mu}(a/(pqt)^{1/2}t; t^2; p, q)}{\Delta_{\mu^2}(a/(pqt)^{1/2}; t; p, q)} \left\langle \frac{\lambda}{\mu^2} \right\rangle_{[a, (pqt)^{1/2}]; t; p, q} = \frac{C_{\lambda}^{-}((pqt)^{1/2}; t; p, q) C_{\lambda}^{+}((pqt)^{1/2}a/t; t; p, q)}{C_{2\lambda}^0((pqt)^{1/2}a/t; t^2; p, q)}. \quad (4.18)$$

One similarly has the following skew analogue.

Theorem 4.4. *The following identity holds:*

$$\sum_{\mu} \mathcal{R}_{\mu^2/\lambda}([v_0, \dots, v_{2n-1}]; ta, (pqt)^{1/2}/a; t; p, q) \frac{\Delta_{\mu}(a^2/(pqt)^{1/2}; t^2; p, q)}{\Delta_{\lambda}(a^2 t/(pqt)^{1/2}V; t; p, q)} \\ \propto \sum_{\mu} \mathcal{R}_{\lambda/\mu^2}([v_0, \dots, v_{2n-1}]; a, (pqt)^{1/2}V/ta; t; p, q) \frac{\Delta_{\mu}(a^2/(pqt)^{1/2}V^2; t^2; p, q)}{\Delta_{\mu^2}(a^2 t/(pqt)^{1/2}V^2; t; p, q)}, \quad (4.19)$$

assuming the LHS terminates; the constant of proportionality is independent of λ , and can be obtained by setting $\lambda = 0$.

Proof. One can again proceed by induction on n ; for $n > 1$, a terminating case always has a pair multiplying to t (possibly after adding a pair multiplying to 1) such that the various sums continue to terminate after extracting that pair. One thus reduces to the case $n = 1$; if $v_0 v_1 = t$, this has already been shown, while in general it follows from Theorem 4.5 below. $\square$

Remark. Again, this formally produces Macdonald's skew q, t -Littlewood identity in the limit.

One disappointing aspect of the above identities is the fact that ab is constrained. It appears that this is a necessary constraint if we wish a completely general Littlewood identity, but if we are willing to restrict our attention to binomial coefficients, we can obtain something more general.

Theorem 4.5. *If $b^2 v_0 v_1 v_2 v_3 = p q t a^2$, and the LHS terminates, then*

$$\sum_{\mu} \left\langle \begin{smallmatrix} \mu^2 \\ \lambda \end{smallmatrix} \right\rangle_{[a,b];t;p,q} \frac{\Delta_{\mu}(a/t|v_0, v_1, v_2, v_3; t^2; p, q)}{\Delta_{\lambda}(a/b|v_0, v_1, v_2, v_3; t; p, q)} \propto \sum_{\mu} \left\langle \begin{smallmatrix} \lambda \\ \mu^2 \end{smallmatrix} \right\rangle_{[a/b,b];t;p,q} \frac{\Delta_{\mu}(a/tb^2|v_0, v_1, v_2, v_3; t^2; p, q)}{\Delta_{\mu^2}(a/b^2|v_0, v_1, v_2, v_3; t; p, q)} \quad (4.20)$$

where the constant of proportionality is independent of λ . The termination condition on the LHS is that

$$t^{2\mathbb{N}} \cap \{v_0, v_1, v_2, v_3, b, b/t\} \neq \emptyset \quad (4.21)$$

$$p^{-\mathbb{N}} q^{-\mathbb{N}} \cap \{v_0, v_1, v_2, v_3, b, b/t\} \neq \emptyset, \quad (4.22)$$

with corresponding constraints on λ if a v_r is used for termination.

Proof. If we write

$$\begin{aligned} \Delta_{\mu}^0(a/t|v_3; t^2; p, q) \left\langle \begin{smallmatrix} \mu^2 \\ \lambda \end{smallmatrix} \right\rangle_{[a,b];t;p,q} &= \Delta_{\mu}^0(a/t|bv_3/t, bv_3/t^2, pqa/v_3; t^2; p, q) \\ &= \sum_{\nu} \left\langle \begin{smallmatrix} \mu^2 \\ \nu \end{smallmatrix} \right\rangle_{[a,t];t;p,q} \left\langle \begin{smallmatrix} \nu \\ \lambda \end{smallmatrix} \right\rangle_{[a/t,b/t](v_3/t, pqa/bv_3);t;p,q} \end{aligned} \quad (4.23)$$

(where v_3 is not used for termination), and apply Lemma 4.1, we reduce to the case with

$$(a, b, v_3) \mapsto (a/t^2, b/t, v_3/t); \quad (4.24)$$

thus by induction, we obtain every case with $b \in t^{\mathbb{N}}$, and the general result by analytic continuation. $\square$

Remark. When $\lambda = 0$, the right-hand sum becomes 1, while the left-hand side becomes

$$\sum_{\mu} \Delta_{\mu}(a/t|v_0, v_1, v_2, v_3, b, b/t; t^2; p, q), \quad (4.25)$$

which can be evaluated, thus determining the normalization.

We observed above that Corollary 3.8 above can be interpreted as giving the inner product of two interpolation functions, and is in particular a special case of a more general integral identity. The same applies to Theorem 4.5. The basic observation is that the sequence of points

$$t^{2n-i}(p, q)^{(\mu^2)_i} a, \quad 1 \leq i \leq 2 \quad (4.26)$$

that arises when evaluating an interpolation function at μ^2 , can also be expressed in the form

$$t^{\pm 1/2} (t^2)^{n-i} (p, q)^{\mu_i} t^{1/2} a, \quad 1 \leq i \leq n. \quad (4.27)$$

This gives rise to the following result, where we recall that

$$\langle f(z_1, \dots, z_n) \rangle_{t_0, t_1, t_2, t_3, u_0, u_1; t; p, q}^{(n)} \quad (4.28)$$

is the normalized linear functional associated to the n -dimensional elliptic Selberg integral.

Theorem 4.6. *For any partition pair λ , and generic parameters satisfying the balancing condition*

$$t^{4n-2}t_0t_1t_2t_3u_0^2 = pq, \quad (4.29)$$

one has

$$\begin{aligned} & \langle \mathcal{R}_\lambda^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, u_0; t; p, q) \rangle_{t^{1/2}t_0, t^{1/2}t_1, t^{1/2}t_2, t^{1/2}t_3, t^{\pm 1/2}u_0; t^2; p, q}^{(n)} \\ &= \Delta_\lambda^0(t^{2n-1}t_0/u_0 | t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3; t; p, q) \\ & \quad \sum_{\mu} \left\langle \frac{\lambda}{\mu^2} \right\rangle_{[t^{2n-1}t_0/u_0, t^{2n-1}t_0u_0]; t; p, q} \frac{\Delta_\mu(1/tu_0^2 | t^{2n}, t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3; t^2; p, q)}{\Delta_{\mu^2}(1/u_0^2 | t^{2n}, t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3; t; p, q)} \end{aligned} \quad (4.30)$$

Proof. If $t^{2n-1}t_0t_1 = p^{-l}q^{-m}$, so that the integral reduces to a sum, the identity is a case of Theorem 4.5. But the left-hand side can be computed by expanding

$$\mathcal{R}_\lambda^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, u_0; t; p, q) \quad (4.31)$$

as a linear combination of products of a function

$$\tilde{\mathcal{R}}_\mu^{(n)}(\dots, z_i, \dots; t^{1/2}t_0 : t^{1/2}t_1, t^{1/2}t_2, t^{1/2}t_3; t^{1/2}u_0, t^{-1/2}u_0; t^2; p, q) \quad (4.32)$$

and a function

$$\tilde{\mathcal{R}}_\nu^{(n)}(\dots, z_i, \dots; t^{1/2}t_0 : t^{1/2}t_1, t^{1/2}t_2, t^{1/2}t_3; t^{-1/2}u_0, t^{1/2}u_0; t^2; p, q). \quad (4.33)$$

(This is not to say that this expansion can be done explicitly; it suffices that such an expansion exists, which follows from the fact that all allowed poles are covered.) In particular, it follows that the left-hand side is the product of p - and q -theta functions, as is the right-hand side, so we may analytically continue to obtain the desired result. $\square$

If we multiply both sides by

$$\Delta_\lambda^0(t^{2n-1}t_0/u_0 | t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3; t; p, q)^{-1} \left\langle \frac{\kappa}{\lambda} \right\rangle_{[1/u_0^2, 1/t^{2n-1}t_0u_0]; t; p, q} \quad (4.34)$$

and sum over λ , the right-hand sum collapses to a delta function, and thus vanishes unless $\kappa = \mu^2$ for some μ . The effect on the left-hand side is to produce a biorthogonal function, and we thus obtain the following vanishing identity.

Corollary 4.7. *For generic parameters such that $t^{4n-2}t_0t_1t_2t_3u_0^2 = pq$, the integral*

$$\langle \tilde{\mathcal{R}}_\lambda^{(n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0 : t_1, t_2, t_3; u_0, u_0; t; p, q) \rangle_{t^{1/2}t_0, t^{1/2}t_1, t^{1/2}t_2, t^{1/2}t_3, t^{\pm 1/2}u_0; t^2; p, q}^{(n)} \quad (4.35)$$

vanishes unless $\lambda = \mu^2$ for some partition pair μ , in which case it equals

$$\frac{\Delta_\mu(1/tu_0^2 | t^{2n}, t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3, 1/t^{2n-1}t_0u_0, 1/t^{2n}t_0u_0; t^2; p, q)}{\Delta_{\mu^2}(1/u_0^2 | t^{2n}, t^{2n-1}t_0t_1, t^{2n-1}t_0t_2, t^{2n-1}t_0t_3, 1/t^{2n-1}t_0u_0, 1/t^{2n-1}t_0u_0; t; p, q)}. \quad (4.36)$$

Remark. In fact, although we have referred to the functions above as “biorthogonal” functions, since $u_0 = u_1$, they in fact form an orthogonal basis of the appropriate space of functions.

If we fix t_0, t_1, t_2, t_3 and let $p \rightarrow 0$ (solving for u_0 via the balancing condition, so that $u_0 \sim \sqrt{p}$), the biorthogonal functions converge to Koornwinder polynomials, and the density converges to a (different) Koornwinder density. The result is one of the vanishing integrals of [9], *together* with the nonzero values (which were not accessible to the methods used there).

We furthermore conjecture that Theorem 4.6 extends to the following transformation (much as Corollary 3.8 extends to Theorem 9.7 of [8]).

Conjecture 2. *For generic parameters such that $t^{4n-4}t_0^2u_0^2v_0v_1v_2v_3 = p^2q^2$, one has*

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, u_0; t; p, q) \Delta^{(n)}(; t^{\pm 1/2}t_0, t^{\pm 1/2}u_0, v_0, v_1, v_2, v_3; t^2; p, q) \\ &= \prod_{0 \leq r \leq 3} \Delta_{\lambda}^0(t^{2n-1}t_0/u_0 | t^{2n-3/2}t_0v_r; t; p, q) \prod_{1 \leq i \leq 2n} \Gamma_{p,q}(t^{2n-1/2-i}t_0v_r, t^{2n-1/2-i}u_0v_r) \\ & \quad \int \mathcal{R}_{\lambda}^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, u_0; t; p, q) \Delta^{(n)}(; t^{\pm 1/2}t_0, t^{\pm 1/2}u_0, v'_0, v'_1, v'_2, v'_3; t^2; p, q), \end{aligned} \quad (4.37)$$

where $v'_r = pq/t^{2n-2}t_0u_0v_r$.

This is accessible in a number of special cases. When $t^{-1/2}t_0v_0 = pq$, so the left-hand side reduces to the left-hand side of Theorem 4.6, the transformed parameters satisfy $t^{2n-2}t^{1/2}u_0v'_0 = 1$, and thus the right-hand side degenerates to a sum. If one traces through the relevant contour conditions, one finds that the sum is over partitions contained in μ , and one obtains the right-hand side of Theorem 4.6. It follows, then, that any “algebraic” case (i.e., in which both sides can be renormalized to products of p - and q -theta functions) of the conjecture holds.

When $\ell(\lambda) = 1$, the skew interpolation function is independent of t , which implies

$$\mathcal{R}_{\lambda}^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, u_0; t; p, q) = \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i, \dots; t^{1/2}t_0, t^{-1/2}u_0; t^2; p, q) \quad (4.38)$$

when $\ell(\lambda) \leq 1$. Thus in that case, the conjecture becomes a special case of [8, Thm. 9.7]. We also find that the identity for $(l, m)^{2n} + \lambda$ follows trivially from that for λ ; combining these two facts proves the identity when $n = 1$, and then trivially the case $t = 1$. In the case $t^{2n}t_0u_0 = pq$, the interpolation function is of Cauchy type, and thus factors for general λ :

$$\begin{aligned} \mathcal{R}_{\lambda}^{*(2n)}(\dots, t^{\pm 1/2}z_i, \dots; t_0, pq/t^{2n}t_0; t; p, q) &= \mathcal{R}_{\lambda^+}^{*(n)}(\dots, z_i, \dots; t^{1/2}t_0, pq/t^{2n+1/2}t_0; t^2; p, q) \\ & \quad \mathcal{R}_{\lambda^-}^{*(n)}(\dots, z_i, \dots; t^{-1/2}t_0, pq/t^{2n-1/2}t_0; t^2; p, q), \end{aligned} \quad (4.39)$$

and again the identity reduces to the transformation of [8, Thm. 9.7]. Finally, the case $t = q$ (and, by symmetry, $t = p$) can be dealt with via the observation that since the interpolation functions can be expressed as a ratio of determinants, one can express the integrands as products of two determinants (one of half the size), with precisely the correct structure for the integrals to become pfaffians of univariate integrals transforming via the $n = 1$ case.

Of course, the usual Littlewood identity also comes in a dual form, and the same applies at the elliptic level. Since duality breaks the symmetry between p and q , it in particular does not apply at the level of partition pairs. However, we do obtain the following, purely p -elliptic, identity.

Corollary 4.8. *One has the identity*

$$\sum_{\mu} \left\langle \begin{matrix} 2\mu \\ \lambda \end{matrix} \right\rangle_{[a,b];q,t;p} \frac{\Delta_{\mu}(a|v_0, v_1, v_2, v_3; q^2, t; p)}{\Delta_{\lambda}(a/b|v_0, v_1, v_2, v_3; q, t; p)} \propto \sum_{\mu} \left\langle \begin{matrix} \lambda \\ 2\mu \end{matrix} \right\rangle_{[a/b,b];q,t;p} \frac{\Delta_{\mu}(a/b^2|v_0, v_1, v_2, v_3; q^2, t; p)}{\Delta_{2\mu}(a/b^2|v_0, v_1, v_2, v_3; q, t; p)}, \quad (4.40)$$

subject to the balancing condition

$$v_0 v_1 v_2 v_3 b^2 = p q t a^2, \quad (4.41)$$

and the termination conditions

$$t^{\mathbb{N}} \cap \{v_0, v_1, v_2, v_3, b, b q\} \neq \emptyset \quad (4.42)$$

$$q^{-2\mathbb{N}} \cap \{v_0, v_1, v_2, v_3, b, b q\} \neq \emptyset, \quad (4.43)$$

with associated conditions on λ . The constant is given by the value for $\lambda = 0$.

Analytically continuing the left-hand side to an integral produces the following dual vanishing integral.

Corollary 4.9. *If $t^{2n-2} t_0 t_1 t_2 t_3 u_0^2 = p q$, then*

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)} (; t_0 : t_1, t_2, t_3; u_0, u_0; q, t; p) \rangle_{t_0, t_1, t_2, t_3, u_0, q u_0; t; p, q^2} \quad (4.44)$$

vanishes unless λ is of the form 2μ , when the integral is

$$\frac{\Delta_{\mu}(1/u_0^2 | t^n, t^{n-1} t_0 t_1, t^{n-1} t_0 t_2, t^{n-1} t_0 t_3, 1/t^{n-1} t_0 u_0, q/t^{n-1} t_0 u_0; q^2, t; p)}{\Delta_{2\mu}(1/u_0^2 | t^n, t^{n-1} t_0 t_1, t^{n-1} t_0 t_2, t^{n-1} t_0 t_3, 1/t^{n-1} t_0 u_0, 1/t^{n-1} t_0 u_0; q, t; p)}. \quad (4.45)$$

Again, here, the vanishing corresponds to the fact that evaluation at a partition with respect to q^2, t , is also evaluation at the doubled partition with respect to q, t :

$$(q^2)^{\mu_i} t^{n-i} t_0 = q^{2\mu_i} t^{n-i} t_0. \quad (4.46)$$

This continues to hold even for partition pairs:

$$p^{\lambda_i} (q^2)^{\mu_i} t^{n-i} t_0 = p^{\lambda_i} q^{2\mu_i} t^{n-i} t_0. \quad (4.47)$$

This suggests the conjecture that

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)} (; t_0 : t_1, t_2, t_3; u_0, u_0; t; p, q) \rangle_{t_0, t_1, t_2, t_3, u_0, q u_0; t; p, q^2}^{(n)} \quad (4.48)$$

vanishes unless $\lambda = (1, 2)\mu$ for some partition pair μ , when it equals

$$\frac{\Delta_{\mu}(1/u_0^2 | t^n, t^{n-1} t_0 t_1, t^{n-1} t_0 t_2, t^{n-1} t_0 t_3, 1/t^{n-1} t_0 u_0, q/t^{n-1} t_0 u_0; t; p, q^2)}{\Delta_{(1,2)\mu}(1/u_0^2 | t^n, t^{n-1} t_0 t_1, t^{n-1} t_0 t_2, t^{n-1} t_0 t_3, 1/t^{n-1} t_0 u_0, 1/t^{n-1} t_0 u_0; t; p, q)}. \quad (4.49)$$

Note, however, that this does not correspond to a vanishing result with respect to the other partition.

The transformation analogue of this extended conjecture is apparently the following.

Conjecture 3. If $t^{2n-2}t_0^2u_0^2v_0v_1v_2v_3 = p^2q^2$, then

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(n)}(; t_0, u_0; t; p, q) \Delta^{(n)}(; t_0, qt_0, u_0, qu_0, v_0, v_1, v_2, v_3; t; p, q^2) \\ &= \prod_{0 \leq r \leq 3} \Delta_{\lambda}^0(t^{n-1}t_0/u_0 | t^{n-1}t_0v_r; t; p, q) \prod_{1 \leq i \leq n} \Gamma_{p,q}(t^{n-i}t_0v_r, t^{n-i}u_0v_r) \\ & \int \mathcal{R}_{\lambda}^{*(n)}(; t_0, u_0; t; p, q) \Delta^{(n)}(; t_0, qt_0, u_0, qu_0, v'_0, v'_1, v'_2, v'_3; t; p, q^2), \end{aligned} \quad (4.50)$$

where $v'_r = pq/t^{n-1}t_0u_0v_r$.

Remark. If $\lambda = (l, m)^n$ (so in particular if $n = 1$), or $t^n t_0 u_0 = pq$, this is again a special case of the transformation of [8, §9].

Consider the q -abelian part of this conjecture, which, if we swap p and q , gives

$$\sum_{\mu} \left\langle \frac{\mu}{\lambda} \right\rangle_{[a,b];q,t;p} \frac{\Delta_{\mu}(a|v_0, v_1, v_2, v_3; q, t; p^2)}{\Delta_{\lambda}(a/b|v_0, v_1, v_2, v_3; q, t; p)} \propto \sum_{\mu} \left\langle \frac{\lambda}{\mu} \right\rangle_{[a/b,b];q,t;p} \frac{\Delta_{\mu}(a/b^2|v_0, v_1, v_2, v_3; q, t; p^2)}{\Delta_{\mu}(a/b^2|v_0, v_1, v_2, v_3; q, t; p)}, \quad (4.51)$$

and thus relates p -abelian and p^2 -abelian functions. This transformation is taken to itself by duality, but if we use a modular transformation, we can replace the 2-isogeny

$$z \in \mathbb{C}^*/\langle p^2 \rangle \mapsto z \in \mathbb{C}^*/\langle p \rangle \quad (4.52)$$

with

$$z \in \mathbb{C}^*/\langle p \rangle \mapsto z^2 \in \mathbb{C}^*/\langle p^2 \rangle. \quad (4.53)$$

This then gives rise to the following conjecture, upon lifting back to an integral transformation.

Conjecture 4. If $t^{n-1}t_0^2u_0^2v_0v_1v_2v_3 = pq$, then

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i^2, \dots; t_0^2, u_0^2; t; p, q) \Delta^{(n)}(; t_0, -t_0, u_0, -u_0, v_0, v_1, v_2, v_3; t^{1/2}; p^{1/2}, q^{1/2}) \\ &= \prod_{0 \leq r \leq 3} \Delta_{\lambda}^0(t^{n-1}t_0^2/u_0^2 | t^{n-1}t_0^2v_r^2; t; p, q) \prod_{0 \leq i < n} \Gamma_{p,q}(t^i t_0^2 v_r^2, t^i u_0^2 v_r^2) \\ & \int \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i^2, \dots; t_0^2, u_0^2; t; p, q) \Delta^{(n)}(; t_0, -t_0, u_0, -u_0, v'_0, v'_1, v'_2, v'_3; t^{1/2}; p^{1/2}, q^{1/2}), \end{aligned} \quad (4.54)$$

where $v'_r = p^{1/2}q^{1/2}/t^{(n-1)/2}t_0u_0v_r$.

Again, this holds if $\lambda = (l, m)^n$ or $t^{n/2}t_0u_0 = p^{1/2}q^{1/2}$.

Although it does not have an associated vanishing result per se, there is an associated analogue of Corollary 4.7, namely the conjecture that if $t^{n-1}t_0t_1t_2t_3u_0^2 = -p^{1/2}q^{1/2}$, then

$$\begin{aligned} & \langle \tilde{R}_{\lambda}^{(n)}(\dots, z_i^2, \dots; t_0^2t_1^2, t_2^2, t_3^2; u_0^2, u_0^2; t; p, q) \rangle_{t_0, t_1, t_2, t_3, u_0, -u_0; t^{1/2}; p^{1/2}, q^{1/2}}^{(n)} \\ &= \frac{\Delta_{\lambda}(1/u_0^2 | t^{n/2}, t^{(n-1)/2}t_0t_1, t^{(n-1)/2}t_0t_2, t^{(n-1)/2}t_0t_3, 1/t^{(n-1)/2}t_0u_0, -1/t^{(n-1)/2}t_0u_0; t^{1/2}; p^{1/2}, q^{1/2})}{\Delta_{\lambda}(1/u_0^4 | t^n, t^{n-1}t_0^2t_1^2, t^{n-1}t_0^2t_2^2, t^{n-1}t_0^2t_3^2, 1/t^{n-1}t_0^2u_0^2, 1/t^{n-1}t_0^2u_0^2; t; p, q)}. \end{aligned} \quad (4.55)$$

5 Quadratic transformations

In [9], there were several other integrals that vanished unless a given partition (or its conjugate) was of the form μ^2 . If we restate the integrals in terms of interpolation polynomials rather than Koornwinder polynomials, the right-hand side becomes a sum over binomial coefficients $\binom{\lambda}{\mu^2}$, multiplied by the nonzero values, suggesting that it should be a special case of the right-hand side of Theorem 4.6. For most of the results of [9], the nonzero values were not established, but in the case of Theorem 4.8 op. cit., they are known, and one can thus use Theorem 4.6 as a guide to formulating the following conjecture.

Conjecture 5. *For generic parameters satisfying $t^{4n-1}t_0^2t_1^2u_0^2 = pq$, the integral*

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(2n)}(\dots, \pm\sqrt{-z_i}, \dots; t_0\sqrt{-1}, -t_0\sqrt{-1}, t_1\sqrt{-1}, -t_1\sqrt{-1}, u_0\sqrt{-1}, -tu_0\sqrt{-1}; t; p, q) \rangle_{t_0^2, t_1^2, u_0^2, t, pt, qt; t^2; p^2, q^2}^{(n)} \quad (5.1)$$

vanishes unless λ is of the form μ^2 , in which case the integral is

$$\frac{\Delta_{\mu}(1/t^2u_0^2|t^{2n}, t^{2n-1}t_0^2, 1/t^{2n-1}t_0u_0, 1/t^{2n}t_0u_0; t^2; p, q)}{\Delta_{\mu^2}(1/tu_0^2|t^{2n}, t^{2n-1}t_0^2, 1/t^{2n-1}t_0u_0, 1/t^{2n}t_0u_0; t; p, q)}. \quad (5.2)$$

If one solves for u_0 in the balancing condition then lets $p \rightarrow 0$, this of course becomes Theorem 4.8 of [9] (and agrees with the nonzero values computed in [6]); one also notes that the conjecture is consistent under negating t_0 or swapping t_0 and t_1 .

There is an alternate formulation of this conjecture as an identity of “hypergeometric” sums. The key observation is that the Cauchy-type interpolation function satisfies the transformation

$$\mathcal{R}_{\lambda}^{*(2n)}(\dots, \pm\sqrt{-z_i}, \dots; pq/t^{2n}\sqrt{-1}u_0, \sqrt{-1}u_0; t; p, q) = \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i, \dots; p^2q^2/t^{2n}u_0^2, u_0^2; t^2; p^2, q^2), \quad (5.3)$$

which follows immediately from the product formula for such functions. (Recall also that, as remarked after Theorem 2.5, we may freely extend the right-hand side to the case $\ell(\lambda) > n$, without invalidating our further computations.)

Thus if the conjecture holds, one can compute the integral

$$\langle \mathcal{R}_{\lambda}^{*(n)}(; p^2q^2/t^{2n}u_0^2, u_0^2; t^2; p^2, q^2) \rangle_{t_0^2, t_1^2, u_0^2, t, pt, qt; t^2; p^2, q^2}^{(n)} \quad (5.4)$$

in two different ways: either by expanding

$$\mathcal{R}_{\lambda}^{*(n)}(; p^2q^2/t^{2n}u_0^2, u_0^2; t^2; p^2, q^2) = \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[p^2q^2/t^2u_0^4, p^2q^2/t^{2n}t_0^2u_0^2](p^2q^2t_0^2/t^2u_0^2); t^2; p^2, q^2} \mathcal{R}_{\mu}^{*(n)}(; t_0^2, u_0^2; t^2; p^2, q^2) \quad (5.5)$$

and applying the elliptic analogue of Kadell’s lemma [8, Cor. 9.3], or by expanding

$$\begin{aligned} \mathcal{R}_{\lambda}^{*(2n)}(; pq/t^{2n}\sqrt{-1}u_0, \sqrt{-1}u_0; t; p, q) \\ = \sum_{\mu} \left\langle \begin{matrix} \lambda \\ \mu \end{matrix} \right\rangle_{[-pq/tu_0^2, -pq/t^{2n}t_0u_0](pqt_0/tu_0); t; p, q} \mathcal{R}_{\mu}^{*(2n)}(; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \end{aligned} \quad (5.6)$$

and applying the conjecture in the form

$$\begin{aligned} & \langle R_{\mu}^{*(2n)}(\dots, \pm\sqrt{-z_i}, \dots; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \rangle_{t_0^2, t_1^2, u_0^2, t, pt, qt; t^2, p^2, q^2}^{(2n)} \\ &= \Delta_{\mu}^0(t^{2n-1}t_0/u_0|t^{2n-1}t_0^2, \pm t^{2n-1}t_0t_1; t; p, q) \sum_{\nu} \left\langle \frac{\mu}{\nu^2} \right\rangle_{[t^{2n-1}t_0/u_0, t^{2n}t_0u_0]; t; p, q} \frac{\Delta_{\nu}(1/t^2u_0^2|t^{2n}, t^{2n-1}t_0^2; t^2; p, q)}{\Delta_{\nu^2}(1/tu_0^2|t^{2n}, t^{2n-1}t_0^2; t; p, q)}. \end{aligned} \quad (5.7)$$

One thus finds that the conjecture implies

$$\begin{aligned} & \sum_{\mu} \left\langle \frac{\lambda}{\mu} \right\rangle_{[p^2q^2/t^2u_0^4, p^2q^2/t^{2n}t_0^2u_0^2](p^2q^2t_0^2/t^2u_0^2); t^2; p^2, q^2} \Delta_{\mu}^0(t^{2n-2}t_0^2/u_0^2|t^{2n-2}t_0^2t_1^2, t^{2n-1}t_0^2, t^{2n-1}pt_0^2, t^{2n-1}qt_0^2; t^2, p^2, q^2) \\ &= \sum_{\mu, \nu} \Delta_{\mu}^0(t^{2n-1}t_0/u_0|t^{2n-1}t_0^2, \pm t^{2n-1}t_0t_1; t; p, q) \frac{\Delta_{\nu}(1/t^2u_0^2|t^{2n}, t^{2n-1}t_0^2; t^2; p, q)}{\Delta_{\nu^2}(1/tu_0^2|t^{2n}, t^{2n-1}t_0^2; t; p, q)} \\ & \quad \left\langle \frac{\lambda}{\mu} \right\rangle_{[-pq/tu_0^2, -pq/t^{2n}t_0u_0](pqt_0/tu_0); t; p, q} \left\langle \frac{\mu}{\nu^2} \right\rangle_{[t^{2n-1}t_0/u_0, t^{2n}t_0u_0]; t; p, q}. \end{aligned} \quad (5.8)$$

In fact, this is equivalent to the conjecture, since one can equally well expand the biorthogonal functions in Cauchy-type interpolation functions. Note also that n enters here only via t^{2n} , and thus one can analytically continue in this extra parameter.

Proposition 5.1. *Conjecture 5 holds whenever $\ell(\lambda) \leq 1$.*

Proof. By triangularity, it suffices to prove (5.8) when $\ell(\lambda) \leq 1$. On the right-hand side, this forces $\ell(\mu) \leq 1$, so $\ell(\nu^2) \leq 1$, and thus $\nu = 0$, making the double sum on the right collapse to a single sum. The resulting identity of univariate elliptic hypergeometric sums is a special case of a known quadratic transformation [14] (the discrete version of Proposition 5.9 below). $\square$

The integral analogue of Conjecture 5 appears to be the following. The label “ $(-1, t^{-1/2})$ ” and similar labels below will be explained at the end of the section.

Conjecture 6. $(-1, t^{-1/2})$. *For otherwise generic parameters satisfying $t^{2n}t_0t_1t_2u_0 = -pq$, one has*

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(2n)}(\dots, \pm\sqrt{-z_i}, \dots; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \Delta^{(n)}(\dots, z_i, \dots; t_0^2, t_1^2, t_2^2, u_0^2, t, pt, qt, pqt; t^2; p^2, q^2) \\ &= \frac{\Delta_{\lambda}^0(t^{2n-1}t_0/u_0| - t^{2n-1}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{2n-1}t_0/u_0|t^{2n}t_0t_1; t; p, q)} \prod_{0 \leq i < 2n} \prod_{0 \leq r < s < 3} \frac{\Gamma_{p,q}(-t^i t_r t_s)}{\Gamma_{p,q}(t^{i+1} t_r t_s)} \\ & \quad \int \mathcal{R}_{\lambda}^{*(2n)}(\dots, t^{\pm 1/2} z_i, \dots; t^{1/2} t_0, t^{1/2} u_0; t; p, q) \Delta^{(n)}(\dots, z_i, \dots; t_0, tt_0, t_1, tt_1, t_2, tt_2, u_0, tu_0; t^2; p, q). \end{aligned} \quad (5.9)$$

Remark. Note that the Δ^0 factor is symmetric under swapping t_1 and t_2 , since the balancing condition makes $(-t^{2n-1}t_0t_1)(t^{2n}t_0t_2) = pqt^{2n-1}t_0/u_0$.

If we use the connection coefficient formula to expand the interpolation functions on each side in terms of the corresponding functions with t_0 replaced by t_1 , the result is a linear combination of instances of the conjecture with t_0 and t_1 swapped. The conjecture is similarly consistent under $\lambda \mapsto (l, m)^{2n} + \lambda$, and (combining the

two) under the Pieri identity. In particular, the case $t_2 = p^l q^m \sqrt{pq/t}$ reduces to the case $t_2 = \sqrt{pq/t}$, which in turn via Theorem 4.6 reduces to Conjecture 5. This, in fact, was how the above conjecture was formulated, by analytically continuing the result of applying the Pieri identity. The fact that the resulting transformation is symmetric in t_1, t_2 is a reassuring consistency; the fact that the right-hand side appears not to be symmetrical under $t_1, t_2 \mapsto -t_1, -t_2$ is less reassuring, but in fact a special case of Conjecture 2 would restore this symmetry.

Proposition 5.2. *Conjecture 6 holds when $n = 1$.*

Proof. Note that this is a nontrivial claim even when $\lambda = 0$, as the two integrands involve different values of p and q . However, we observe that in general the case of the conjecture with $\lambda = (l, m)^{2n} + \mu$ reduces to the case with $\lambda = \mu$, so for $n = 1$, it suffices to consider the case $\ell(\lambda) \leq 1$. But then the integral representation of [8] implies the following expression.

$$\begin{aligned} & \mathcal{R}_{(l,m)}^{*(2)}(\pm\sqrt{-z}; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \\ &= \frac{\Gamma_{p^2, q^2}(t^2 u_0^2 z^{\pm 1})}{\Gamma_{p^2, q^2}(u_0^2 z^{\pm 1}, tz^{\pm 1}, ptz^{\pm 1}, qtz^{\pm 1}, pqtz^{\pm 1})} \times \frac{(p; p)(q; q)\Gamma_{p,q}(t^2)}{2\Gamma_{p^2, q^2}(t^2)^2} \\ & \int_{C'} \mathcal{R}_{(l,m)}^{*(1)}(y; t^{1/2}t_0\sqrt{-1}, t^{-1/2}u_0\sqrt{-1}; t; p, q) \frac{\Gamma_{p,q}(t^{-1/2}u_0y^{\pm 1}\sqrt{-1})\Gamma_{p^2, q^2}(-tz^{\pm 1}y^{\pm 2})}{\Gamma_{p,q}(t^{3/2}u_0y^{\pm 1}\sqrt{-1})\Gamma_{p,q}(y^{\pm 2})} \frac{dy}{2\pi\sqrt{-1}y}. \end{aligned} \quad (5.10)$$

If we substitute this in and exchange order of integration, the integral over z becomes an instance of the order 0 elliptic beta integral (the z -dependent factor above cancels five parameters then adds three, making a final total of six), so can be explicitly evaluated. We thus conclude

$$\begin{aligned} & \frac{(p^2; p^2)(q^2; q^2)\Gamma_{p^2, q^2}(t^2)}{(p; p)(q; q)\Gamma_{p,q}(t^2)} \int \mathcal{R}_{(l,m)}^{*(2)}(\pm\sqrt{-z}; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \Delta^{(1)}(z; t_0^2, t_1^2, t_2^2, u_0^2, t, pt, qt, pqt; t^2, p^2, q^2) \\ &= \prod_{0 \leq r < s < 2} \Gamma_{p^2, q^2}(t_r^2 t_s^2) \prod_{0 \leq r < 2} \Gamma_{p^2, q^2}(t^2 u_0^2 t_r^2) \\ & \int \mathcal{R}_{(l,m)}^{*(1)}(y; t^{1/2}t_0\sqrt{-1}, t^{-1/2}u_0\sqrt{-1}; t; p, q) \\ & \Delta^{(1)}(y; \pm t^{1/2}t_0\sqrt{-1}, \pm t^{1/2}t_1\sqrt{-1}, \pm t^{1/2}t_2\sqrt{-1}, t^{-1/2}u_0\sqrt{-1}, -t^{3/2}u_0\sqrt{-1}; t; p, q). \end{aligned} \quad (5.11)$$

On the right-hand side, we have

$$\mathcal{R}_{(l,m)}^{*(2)}(t^{\pm 1/2}z; t^{1/2}t_0, t^{1/2}u_0; t; p, q) = \mathcal{R}_{(l,m)}^{*(1)}(z; tt_0, u_0; t^2; p, q), \quad (5.12)$$

and thus the integral on the right-hand side is

$$\int \mathcal{R}_{(l,m)}^{*(1)}(z; tt_0, u_0; t^2; p, q) \Delta^{(1)}(z; t_0, tt_0, t_1, tt_1, t_2, tt_2, u_0, tu_0; t^2; p, q) \quad (5.13)$$

But the general univariate interpolation function and the general univariate density are independent of t , so we may replace t^2 by t above. The desired special case of Conjecture 6 then follows as a special case of [8, Cor. 9.11]. $\square$

This immediately implies that Conjecture 6 holds when $t = 1$. Another special case arises when $t = q$ (or, symmetrically, $t = p$), in which case both integrands can be expressed as the product of a determinant of size $2n$ (coming from the p -elliptic half of the interpolation function and the cross terms) and the symmetrization of a product (coming from the q -elliptic half of the interpolation function). Thus both integrals are sums of $(2n)!$ pfaffians, and each entry of each pfaffian transforms via the $n = 1$ instance.

We also have an additional special case when $\lambda = 0$.

Proposition 5.3. *Conjecture 6 holds for $\lambda = 0$, $t \in \{p^{1/2}, q^{1/2}\}$.*

Proof. When $t = q^{1/2}$, two parameters cancel in the left-hand side, allowing it to be evaluated, while the right-hand side can be evaluated by observing that its integrand is equal to an elliptic Selberg integrand of order 0 with $q \mapsto q^{1/2}$; the result follows upon simplifying the gamma factors. $\square$

The above evaluation of the right-hand side generalizes to a transformation, again by observing that both sides have the same integrand.

Proposition 5.4. *For any odd integer $m > 0$,*

$$II_n^{(m)}(u_0, qu_0, \dots, u_{m+2}, qu_{m+2}; q^2; p, q^2) = II_n^{((m-1)/2)}(u_0, \dots, u_{m+2}, \pm q^{1/2}; q; p, q), \quad (5.14)$$

subject to the balancing condition $q^{2n-1} \prod_{0 \leq r < m+3} u_r = -(pq)^{(m+1)/2}$.

Remark. When $m = 1$, this is the aforementioned evaluation. One can relax the condition that m is odd by taking $u_{m+2} = \pm p^{1/2} q^{1/2}$, thus causing a pair of parameters to cancel on the left, but not the right; similarly, one can change the sign of the balancing condition at the cost of increasing the order on the right.

A similar argument gives the following, univariate only, transformation.

Proposition 5.5. *For any even integer $m \geq 0$,*

$$I^{(m)}(\pm \sqrt{u_0}, \dots, \pm \sqrt{u_{m+2}}; p, q) = 2\Gamma_{p,q}(-1)I^{(m/2)}(u_0, \dots, u_{m+2}, -1, -q, -p; p^2, q^2), \quad (5.15)$$

subject to the balancing condition $\prod_{0 \leq r < m+3} u_r = -(pq)^{m+1}$.

Since Conjecture 5 involves a choice of 4-torsion point on the elliptic curves (namely $\sqrt{-1}$), it has an equivariant form under modular transformation. This should then extend back to general partition pairs, although we have rather less guidance in this case. Luckily, the argument for $n = 1$ carries over with little change. The resulting integral breaks symmetry between u_1, u_2 , but this can be restored by adding an additional parameter, as follows.

Conjecture 7. *($p^{1/2}, t^{-1/2}$) For otherwise generic parameters satisfying $t^{2n} t_0 t_1 t_2 u_0 = p^{1/2} q$, one has*

$$\begin{aligned} & \int \mathcal{R}_\lambda^{*(2n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots; p^{1/4} t_0, p^{1/4} u_0; t; p, q) \Delta^{(n)}(; t_0, t_1, t_2, u_0, \pm \sqrt{t}, \pm \sqrt{qt}, p^{1/2} v, p^{1/2} q/v; t; p^{1/2}, q) \\ &= \frac{\Delta_\lambda^0(t^{2n-1} t_0 / u_0 | t^{2n-1} p^{1/2} t_0 t_1; t; p, q)}{\Delta_\lambda^0(t^{2n-1} t_0 / u_0 | t^{2n} t_0 t_1; t; p, q)} \prod_{\substack{0 \leq i < 2n \\ 0 \leq r < s < 3}} \frac{\Gamma_{p,q}(t^i p^{1/2} t_r t_s)}{\Gamma_{p,q}(t^{i+1} t_r t_s)} \\ & \int \mathcal{R}_\lambda^{*(2n)}(\dots, t^{1/2} z_i^{\pm 1}, \dots; t^{1/2} t_0, t^{1/2} u_0; t; p, q) \Delta^{(n)}(; t_0, t t_0, t_1, t t_1, t_2, t t_2, u_0, t u_0, p v, p q/v; t^2; p, q). \end{aligned} \quad (5.16)$$

Remark 1. The additional parameter has the effect of multiplying each integrand by

$$\prod_{1 \leq i \leq n} \theta_q(v z_i^{\pm 1}). \quad (5.17)$$

Since these functions span the $(n + 1)$ -dimensional space of BC_n -symmetric q -theta functions of degree 1, one may replace this factor by an arbitrary such function without affecting the validity of the conjecture. In particular, for $n = 1$, it suffices to verify the conjecture for two values of v , say $v = t_1$, $v = t_2$, which eliminates the extra parameter, and allows the previous argument to apply. The cases $t = 1$, $t = p$, $t = q$ follow as before.

Remark 2. Again, this (and the remainder of the conjectures we will formulate along these lines) is consistent with respect to connection coefficients, $\lambda \mapsto (l, m)^{2n} + \lambda$, and the Pieri identity, regardless of the additional parameter. Another important consistency condition is that if we take $v = t_2$ then multiply t_2 by $p^{-1/2}$, the left-hand side is again symmetric in t_1 and t_2 , while an application of Conjecture 2 exhibits the corresponding symmetry on the right-hand side.

The corresponding vanishing conjecture (obtained by taking $v = t_2$ to eliminate the extra parameter, then $t_2 = q^{1/2}t^{-1/2}$ or $t_1 = p^{1/2}q^{1/2}t^{-1/2}$, then applying Theorem 4.6) is as follows.

Conjecture 8. *For generic parameters satisfying $t^{2n-1/2}t_0t_1u_0 = p^{1/2}q^{1/2}$, the integral*

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(2n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots; p^{1/4}t_0, p^{-1/4}t_0, p^{1/4}t_1, p^{-1/4}t_1; p^{1/4}u_0, p^{-1/4}tu_0; t; p, q) \rangle_{t_0, t_1, u_0, t^{1/2}, -t^{1/2}, -q^{1/2}t^{1/2}; t; p^{1/2}, q}^{(n)} \quad (5.18)$$

vanishes unless λ is of the form μ^2 , in which case the integral is

$$\frac{\Delta_{\mu}(1/t^2u_0^2|t^{2n}, t^{2n-1}t_0^2, 1/t^{2n-1}t_0u_0, 1/t^{2n}t_0u_0; t^2; p, q)}{\Delta_{\mu^2}(1/tu_0^2|t^{2n}, t^{2n-1}t_0^2, 1/t^{2n-1}t_0u_0, 1/t^{2n}t_0u_0; t; p, q)}. \quad (5.19)$$

Remark. Note that the nonzero values are the same as in Conjecture 5.

This would imply Conjecture 5 via a modular transformation, as discussed above, but the q -elliptic half of the identity would be new. In the limit $q \rightarrow 0$, t_0, t_1 fixed of that q -elliptic identity, the biorthogonal function becomes a Koornwinder polynomials, and one obtains the vanishing identity given as Theorem 4.10 of [9], together with a conjecture for the nonzero values. The case $t_0 = q^{1/2}t^{1/2}$, $t_1 \mapsto p^{1/4}a$ is also of interest, as in that case the biorthogonal function becomes a suitably normalized interpolation function. One can then take the limit $p \rightarrow 0$ with a fixed, in which limit the integral becomes

$$\int \frac{P_{\lambda}(\dots, z_i^{\pm 1}, \dots; q, t)}{P_{\lambda}(q^{-1/2}t^{1/2-2n}, \dots, q^{-1/2}t^{-1/2}; q, t)} \prod_{1 \leq i < j \leq n} \frac{(z_i^{\pm 1}z_j^{\pm 1}; q)}{(tz_i^{\pm 1}z_j^{\pm 1}; q)} \prod_{1 \leq i \leq n} \frac{(z_i^{\pm 2}; q)}{(tz_i^{\pm 2}; q)} \frac{dz_i}{2\pi\sqrt{-1}z_i}. \quad (5.20)$$

Apart from the change in normalization of the Macdonald polynomial, this integral is that of Theorem 4.1 of [9], and therefore vanishes unless $\lambda = \mu^2$. Moreover, the corresponding nonzero values (known in this case) agree with those obtained by degenerating Conjecture 8.

Again, there is a sum version, this time based on the identity

$$\mathcal{R}_{\lambda}^{*(2n)}(\dots p^{1/4}z_i^{\pm 1} \dots; p^{3/4}q/t^{2n}u_0, p^{1/4}u_0; t; p, q) = \mathcal{R}_{(2,1)\lambda}^{*(n)}(\dots z_i \dots; p^{1/2}q/t^n u_0, u_0; t; p^{1/2}, q) \quad (5.21)$$

of Cauchy-type interpolation functions (suitably extended); we omit the details. When $\ell(\lambda) \leq 1$, the sum is a special case of the discrete version of Proposition 5.6 below (which discrete version in turn combines a quadratic transform of Warnaar [15] with the modular transform of the transform of Spiridonov [14] mentioned above).

Of course, the next step is to dualize the above conjectures; however, we see by reference to the known trigonometric cases that some subtleties will arise. The vanishing integral of Theorem 4.8 of [9] (corresponding to Conjecture 5) actually dualizes to a pair of vanishing integrals, while for Theorem 4.1 of [9] (a limit of Conjecture 8), not only are there two dual identities, but each dual identity itself involves a sum of two integrals.

One case is straightforward, namely the “other” dual of Conjecture 7 (i.e., exchange p and q before dualizing). Here, and in the other two cases, we begin by dualizing the algebraic versions of the conjectures, a la (5.8), having first analytically continued in $t^{2n} = T$. After reparametrizing and specializing T appropriately, we can recognize the left-hand side as the integral of a Cauchy-type interpolation function, and thus re-express the dual as a vanishing identity. At that point, one may use the Pieri identity to extend to a large set of cases of an integral transformation.

Conjecture 9. $(t^{-1/2}, q^{1/2})$ Subject to the balancing conditions $t^{n-1}t_0t_1t_2u_0 = pq^2t^{1/2}$, $tv_0v_1 = pq$, one has

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(n)}(; t^{-1/4}t_0, t^{-1/4}u_0; t; p, q) \\ & \Delta^{(n)}(; t^{-1/4}t_0, t^{-1/4}t_1, t^{-1/4}t_2, t^{-1/4}u_0, \pm t^{1/4}, \pm p^{1/2}t^{1/4}, t^{1/4}v_0, t^{1/4}v_1; t^{1/2}; p, q) \\ & = \frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-3/2}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1/q; t; p, q)} \prod_{0 \leq i < n} \prod_{0 \leq r < s < 3} \frac{\Gamma_{p,q}(t^{i-1/2}t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s / q)} \\ & \int \mathcal{R}_{\lambda}^{*(n)}(; q^{-1/2}t_0, q^{-1/2}u_0; t; p, q) \Delta^{(n)}(; q^{\pm 1/2}t_0, q^{\pm 1/2}t_1, q^{\pm 1/2}t_2, q^{\pm 1/2}u_0, q^{1/2}v_0, q^{1/2}v_1; t; p, q^2), \end{aligned} \quad (5.22)$$

Remark 3. Here the extra parameter multiplies the integrands by factors

$$\prod_{0 \leq i < n} \frac{\Gamma_{p,q}(t^{1/4}v_0z_i^{\pm 1})}{\Gamma_{p,q}(t^{3/4}v_0z_i^{\pm 1})} \quad \text{and} \quad \prod_{0 \leq i < n} \frac{\Gamma_{p,q^2}(q^{1/2}v_0z_i^{\pm 1})}{\Gamma_{p,q^2}(q^{1/2}v_0z_i^{\pm 1})}, \quad (5.23)$$

which are not, in fact, theta functions. They do, however, closely resemble the generating function for the q, t -analogues g_k of the complete symmetric functions [4].

Remark. If we eliminate the extra parameters from the normalization (i.e., $\lambda = 0$) cases of this conjecture and Conjecture 7, the resulting quadratic transformations are equivalent up to an E_7 transformation of the right-hand side and reparametrizing.

Proposition 5.6. Conjecture 9 holds when $n = 1$.

Proof. Since $\ell(\lambda) \leq 1$, we immediately reduce to the case $\lambda = 0$, for which we need simply exchange order of integration in the double integral

$$\int \int \Gamma_{p,q}(q^{-1/2}t^{1/4}x^{\pm 1}y^{\pm 1}) \frac{\prod_{0 \leq r < 4} \Gamma_{p,q}(t^{-1/4}u_r x^{\pm 1})}{\Gamma_{p,q}(x^{\pm 2})} \frac{dx}{2\pi\sqrt{-1}x} \frac{\Gamma_{p,q^2}(q^{1/2}v_0y^{\pm 1}, q^{1/2}v_1y^{\pm 1})}{\Gamma_{p,q^2}(y^{\pm 2})} \frac{dy}{2\pi\sqrt{-1}y} \quad (5.24)$$

and simplify the resulting integrands. $\square$

Unfortunately, this case is not sufficient to prove the $t = p$ and $t = q$ cases, as although they can again (and for the later conjectures) be expressed via pfaffians, the entries of the pfaffians include instances with $n = 2$. The univariate case does, however, suffice to prove the case $t = 1$.

The corresponding vanishing conjecture states that

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(; t^{-1/4}t_0, t^{1/4}t_0, t^{-1/4}t_1, t^{1/4}t_1; t^{-1/4}u_0, t^{1/4}u_0/q; t; p, q) \rangle_{t^{-1/4}t_0, t^{-1/4}t_1, t^{-1/4}u_0, t^{1/4}, -t^{1/4}, -p^{1/2}t^{1/4}; t^{1/2}; p, q}^{(n)} \quad (5.25)$$

vanishes unless $\lambda = (1, 2)\mu$, when it equals

$$\frac{\Delta_{\mu}(q/u_0^2|t^n, t^{n-1}t_0^2, 1/t^{n-1}t_0u_0, q/t^{n-1}t_0u_0; t; p, q^2)}{\Delta_{(1,2)\mu}(q/u_0^2|t^n, t^{n-1}t_0^2, 1/t^{n-1}t_0u_0, q/t^{n-1}t_0u_0; t; p, q)}. \quad (5.26)$$

The next simplest case is the dual of Conjecture 6. Here we find that the integral on the left-hand side is half the dimension of that on the right-hand side, which is thus necessarily even. This constraint can be avoided, however, by observing that the corresponding integral of Cauchy-type interpolation functions can again be expressed as such an integral, even when the dimension is apparently a half-integer. One thus obtains the following conjecture.

Conjecture 10. $(-1, q^{1/2})$ For otherwise generic parameters satisfying $t^{n-1}t_0t_1t_2u_0 = -pq^2$, the integral

$$\frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0| -t^{n-1}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0| t^{n-1}t_0t_1/q; t; p, q)} \prod_{0 \leq i < n} \prod_{0 \leq r < s < 3} \frac{\Gamma_{p,q}(-t^i t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s/q)} \int \mathcal{R}_{\lambda}^{*(n)}(\dots z_i \dots; q^{-1/2}t_0, q^{-1/2}u_0; t; p, q) \Delta^{(n)}(; q^{\pm 1/2}t_0, q^{\pm 1/2}t_1, q^{\pm 1/2}t_2, q^{\pm 1/2}u_0; t; p, q^2) \quad (5.27)$$

is equal to

$$\int \mathcal{R}_{\lambda}^{*(n)}(\dots \pm \sqrt{-z_i} \dots; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \Delta^{(n/2)}(; t_0^2, t_1^2, t_2^2, u_0^2, 1, p, t, pt; t^2; p^2, q^2) \quad (5.28)$$

if n is even, and

$$\Gamma_{p^2, q^2}(t_0^2, t_1^2, t_2^2, u_0^2, p, t, pt) \int \mathcal{R}_{\lambda}^{*(n)}(\dots \pm \sqrt{-z_i} \dots, \sqrt{-1}; t_0\sqrt{-1}, u_0\sqrt{-1}; t; p, q) \Delta^{((n-1)/2)}(; t_0^2, t_1^2, t_2^2, u_0^2, t^2, p, t, pt; t^2; p^2, q^2) \quad (5.29)$$

if n is odd.

Proposition 5.7. Conjecture 10 holds when $\lambda = 0$ and $t = q^2$.

Proof. As in Proposition 5.3, one side is an elliptic Selberg integral, while the other can be evaluated using Proposition 5.4. $\square$

In particular, when $n = 1$, both sides are essentially independent of t , and thus the normalization conjecture (i.e., the case $\lambda = 0$) holds in that case as well, which in turn implies the $n = 1$ case of Conjecture 10 in its entirety. Again, however, a proof for $t \in \{p, q\}$ would require the corresponding $n = 2$ cases to be demonstrated. Even the case $t = 1$ is rather subtle in this case, since then the relevant univariate integrals become singular.

The corresponding vanishing conjecture (take $t_2 = p^{l+1/2}q^{m+1}$) reads that the integrals

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, \pm\sqrt{-z_i}, \dots; t_0\sqrt{-1}: -t_0\sqrt{-1}, \pm t_1\sqrt{-1}; u_0\sqrt{-1}, -u_0\sqrt{-1}/q; t; p, q) \rangle_{t_0^2, u_0^2, t_1^2, t^2, t, pt; t^2; p^2, q^2}^{(n/2)} \quad (5.30)$$

and

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, \pm\sqrt{-z_i}, \dots, \sqrt{-1}; t_0\sqrt{-1}: -t_0\sqrt{-1}, \pm t_1\sqrt{-1}; u_0\sqrt{-1}, -u_0\sqrt{-1}/q; t; p, q) \rangle_{t_0^2, u_0^2, t_1^2, t^2, t, pt; t^2; p^2, q^2}^{((n-1)/2)} \quad (5.31)$$

subject to the balancing condition $t^{2n-2}t_0^2t_1^2u_0^2 = pq^2$, vanish unless λ has the form $(1, 2)\mu$, when the value is as in (5.26) above. The $n = 1$ instance of this is a quadratic evaluation formula due to Warnaar [16].

The remaining case is the direct dual of Conjecture 8. If we attempt to proceed as above, we find that the most straightforward version of the half-integer case fails to hold; although taking $t_1 = p^{1/2}q$ reduces to the dual vanishing identity, taking $t_1 = -p^{1/2}q$ makes the right-hand side vanish. If one instead takes a sum of two terms, symmetric under $p^{1/2} \mapsto -p^{1/2}$, this problem disappears. This, of course, corresponds to the fact that the known Macdonald polynomial limit itself involves a sum of two integrals. The corresponding structure for the integer case is then reasonably straightforward to guess. One thus formulates the following conjecture.

Conjecture 11. ($p^{1/2}, q^{1/2}$) *For otherwise generic parameters satisfying $t^{n-1}t_0t_1t_2u_0 = p^{1/2}q^2$, the rescaled integral*

$$\frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}p^{1/2}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1/q; t; p, q)} \prod_{\substack{0 \leq i < n \\ 0 \leq r < s < 3}} \frac{\Gamma_{p,q}(t^i p^{1/2} t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s / q)} \int \mathcal{R}_{\lambda}^{*(n)}(; q^{-1/2}t_0, q^{-1/2}u_0; t; p, q) \Delta^{(n)}(; q^{\pm 1/2}t_0, q^{\pm 1/2}t_1, q^{\pm 1/2}t_2, q^{\pm 1/2}u_0, pqv^{\pm 1}; t; p, q^2) \quad (5.32)$$

admits the following expressions as sums of lower-dimensional integrals. If n is odd, then it equals

$$\begin{aligned} & \Gamma_{p^{1/2}, q}(t_0, t_1, t_2, u_0, -1, \pm t^{1/2}, p^{1/2}q^{1/2}v^{\pm 1}) \\ & \int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, p^{1/4}; p^{1/4}t_0, p^{1/4}u_0; t; p, q) \\ & \Delta^{((n-1)/2)}(; t_0, t_1, t_2, u_0, t, -1, \pm t^{1/2}, p^{1/2}q^{1/2}v^{\pm 1}; t; p^{1/2}, q) \\ & + \Gamma_{p^{1/2}, q}(-t_0, -t_1, -t_2, -u_0, -1, \pm t^{1/2}, -p^{1/2}q^{1/2}v^{\pm 1}) \\ & \int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, -p^{1/4}; p^{1/4}t_0, p^{1/4}u_0; t; p, q) \\ & \Delta^{((n-1)/2)}(; t_0, t_1, t_2, u_0, 1, -t, \pm t^{1/2}, p^{1/2}q^{1/2}v^{\pm 1}; t; p^{1/2}, q), \end{aligned} \quad (5.33)$$

while if n is even, it equals

$$\begin{aligned}
& \int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots; p^{1/4} t_0, p^{1/4} u_0; t; p, q) \\
& \quad \Delta^{(n/2)}(; t_0, t_1, t_2, u_0, \pm 1, \pm t^{1/2}, p^{1/2} q^{1/2} v^{\pm 1}; t; p^{1/2}, q) \\
& + \Gamma_{p, q^2}(t_0^2, t_1^2, t_2^2, u_0^2, t, t, p q v^{\pm 2}) \Gamma_{p^{1/2}, q}(-1, -t) \\
& \int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots, \pm p^{1/4}, p^{1/4} t_0, p^{1/4} u_0; t; p, q) \\
& \quad \Delta^{(n/2-1)}(; t_0, t_1, t_2, u_0, \pm t, \pm t^{1/2}, p^{1/2} q^{1/2} v^{\pm 1}; t; p^{1/2}, q). \tag{5.34}
\end{aligned}$$

Remark. Here the extra parameter multiplies the first integrand by

$$f(z_1, \dots, z_n) = \prod_{1 \leq i \leq n} \theta_{q^2}(q v z_i^{\pm 1}), \tag{5.35}$$

while the integrands on the right are multiplied by

$$f(q^{1/2} z_1^{\pm 1}, \dots, q^{1/2} z_{(n-1)/2}^{\pm 1}, q^{1/2}), \quad f(q^{1/2} z_1^{\pm 1}, \dots, q^{1/2} z_{(n-1)/2}^{\pm 1}, -q^{1/2}), \tag{5.36}$$

$$f(q^{1/2} z_1^{\pm 1}, \dots, q^{1/2} z_{n/2}^{\pm 1}), \quad f(q^{1/2} z_1^{\pm 1}, \dots, q^{1/2} z_{n/2-1}^{\pm 1}, \pm q^{1/2}), \tag{5.37}$$

and this factor accounts for all dependence on v . One could thus just as well replace f by an arbitrary BC_n -symmetric q^2 -theta function of degree 1. From this perspective, each of the above four specializations induces a linear transformation from the space of such theta functions to a corresponding space of BC -symmetric q -theta functions. Moreover, each such transformation is surjective, with kernel one of the two eigenspaces of the operator

$$f(z_1, \dots, z_n) \rightarrow q^{n/2} \left(\prod_{1 \leq i \leq n} z_i \right) f(q z_1, \dots, q z_n). \tag{5.38}$$

Proposition 5.8. *Conjecture 11 holds when $n = 1$.*

Proof. Since $\ell(\lambda) \leq 1$ when $n = 1$, we may as well take $\lambda = 0$. We thus need simply to prove that when $u_0 u_1 u_2 u_3 = p^{1/2} q^2$,

$$\begin{aligned}
& \frac{(p; p)(q^2; q^2)}{2} \int \frac{\prod_{0 \leq r < 4} \Gamma_{p, q^2}(q^{\pm 1/2} u_r z^{\pm 1})}{\Gamma_{p, q^2}(z^{\pm 2})} \frac{\theta_{q^2}(q v z^{\pm 1}) dz}{2\pi \sqrt{-1} z} \\
& = \frac{\Gamma_{p^{1/2}, q}(-1, u_0, u_1, u_2, u_3) \theta_q(q^{1/2} v)}{\prod_{0 \leq r < s < 4} \Gamma_{p, q}(p^{1/2} u_r u_s)} + \frac{\Gamma_{p^{1/2}, q}(-1, -u_0, -u_1, -u_2, -u_3) \theta_q(-q^{1/2} v)}{\prod_{0 \leq r < s < 4} \Gamma_{p, q}(p^{1/2} u_r u_s)} \tag{5.39}
\end{aligned}$$

Each of the three terms is a BC_1 -symmetric q^2 -theta function in v of degree 1, and thus the relation will follow if we check it at any two independent points. By symmetry under $v \mapsto -v$, we may reduce to the case $v = q^{-1/2}$, when the left-hand side can be expressed (via Proposition 5.4) as

$$\frac{(q^2; q^2)}{(q; q)} I^{(0)}(q^{-1/2} u_0, q^{-1/2} u_1, q^{-1/2} u_2, q^{-1/2} u_3, -q^{1/2}, -p^{1/2} q^{1/2}; p, q). \tag{5.40}$$

The proposition follows upon simplifying the resulting product of elliptic gamma functions. $\square$

Remark. This can also be obtained as the limit $t \rightarrow 1/p$ of Proposition 5.6; one finds that the left-hand side of that Proposition violates the contour conditions in two different ways in the limit, and thus becomes a sum of two residues, corresponding to the two terms above.

When $t_1 = -p^{1/2}q$, $v = q^{1/2}/t_2$, one of the two terms vanishes, since $\Gamma(p^{1/2}q; p^{1/2}, q) = 0$, and we thus obtain (after an application of Conjecture 3) the following vanishing conjecture; namely, that when $t^{n-1}t_0t_1u_0 = -p^{1/2}q$, the integral

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots; p^{1/4}t_0: p^{-1/4}t_0, p^{1/4}t_1, p^{-1/4}t_1; p^{1/4}u_0, p^{-1/4}u_0/q; t; p, q) \rangle_{t_0, t_1, u_0, 1, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{(n/2)} \quad (5.41)$$

or

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, p^{1/4}; p^{1/4}t_0: p^{-1/4}t_0, p^{1/4}t_1, p^{-1/4}t_1; p^{1/4}u_0, p^{-1/4}u_0/q; t; p, q) \rangle_{t_0, t_1, u_0, t, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{((n-1)/2)} \quad (5.42)$$

as appropriate, vanishes unless $\lambda = (1, 2)\mu$, when its value is given by (5.26).

Similarly, in the limit $t_0t_1 \rightarrow p^{1/2}q$, the n -dimensional integral degenerates to a (dual) Littlewood-style sum, and an application of connection coefficients gives the conjecture that when $t^{n-1}t_0u_0 = p^{1/2}q$, the integrals

$$\begin{aligned} & \frac{1}{2} \langle \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots; p^{-1/4}t_0, p^{1/4}u_0; t; p, q) \rangle_{t_0, u_0, 1, -1, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{(n/2)} \\ & + \frac{1}{2} \langle \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, \pm p^{1/4}; p^{-1/4}t_0, p^{1/4}u_0; t; p, q) \rangle_{t_0, u_0, t, -t, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{(n/2-1)} \end{aligned} \quad (5.43)$$

and

$$\begin{aligned} & \frac{1}{2} \langle \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, p^{1/4}; p^{-1/4}t_0, p^{1/4}u_0; t; p, q) \rangle_{t_0, u_0, t, -1, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{((n-1)/2)} \\ & + \frac{1}{2} \langle \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4}z_i^{\pm 1}, \dots, -p^{1/4}; p^{-1/4}t_0, p^{1/4}u_0; t; p, q) \rangle_{t_0, u_0, 1, -t, t^{1/2}, -t^{1/2}; t; p^{1/2}, q}^{((n-1)/2)} \end{aligned} \quad (5.44)$$

vanish unless $\lambda = (1, 2)\mu$, when they have value

$$\frac{\Delta_{\mu}(q/u_0^2|t^n, t^{n-1}; t; p, q^2)}{\Delta_{(1,2)\mu}(q/u_0^2|t^n, t^{n-1}; t; p, q)}. \quad (5.45)$$

If we swap p and q above, we find that Conjecture 10 becomes self-dual, while Conjectures 9 and 11 become dual to each other. However, we now have the possibility again of modular transformations. Given the lack of guidance from the trigonometric level, the resulting conjectures are rather more speculative than those above. The overall form of the integrals is fairly straightforward to determine, especially since in each case the normalization without extra parameter reduces via an E_7 transformation to a previously conjectured normalization. The λ -dependent factors are then uniquely determined by the requirement of consistency under the Pieri identity (more precisely, that the obvious argument for consistency should work, as it did in all previous cases).

For Conjecture 10, one obtains the following transform, of which the case $n = 1$ is straightforward.

Conjecture 12. $(p^{1/2}, -1)$ For otherwise generic parameters satisfying $t^{n-1}t_0t_1t_2u_0 = p^{1/2}q$, the integral

$$\frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|p^{1/2}t^{n-1}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1; t; p, q)} \prod_{0 \leq i < n} \prod_{0 \leq r < s < 3} \frac{\Gamma_{p,q}(t^i p^{1/2} t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s)} \int \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i^2, \dots; -t_0, -u_0; t; p, q) \Delta^{(n)}(\pm \sqrt{-t_0}, \pm \sqrt{-t_1}, \pm \sqrt{-t_2}, \pm \sqrt{-u_0}, p^{1/2} q^{1/4} v^{\pm 1}; t^{1/2}; p^{1/2}, q^{1/2}) \quad (5.46)$$

is equal to

$$\int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots; p^{1/4} t_0, p^{1/4} u_0; t; p, q) \Delta^{(n/2)}(\pm t_0, \pm t_1, \pm t_2, \pm u_0, 1, q^{1/2}, t^{1/2}, q^{1/2} t^{1/2}, -p^{1/2} q^{1/2} v^{\pm 2}; t; p^{1/2}, q) \quad (5.47)$$

if n is even, and

$$\Gamma_{p^{1/2}, q}(t_0, t_1, t_2, u_0, q^{1/2}, t^{1/2}, q^{1/2} t^{1/2}, -p^{1/2} v^2, -p^{1/2} q/v^2) \int \mathcal{R}_{\lambda}^{*(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots, p^{1/4}; p^{1/4} t_0, p^{1/4} u_0; t; p, q) \Delta^{((n-1)/2)}(\pm t_0, \pm t_1, \pm t_2, \pm u_0, t, q^{1/2}, t^{1/2}, q^{1/2} t^{1/2}, -p^{1/2} q^{1/2} v^{\pm 2}; t; p^{1/2}, q) \quad (5.48)$$

if n is odd.

Remark 1. Once again, the factor

$$\prod_{1 \leq i \leq n} \Gamma_{p^{1/2}, q^{1/2}}(p^{1/2} q^{1/4} v^{\pm 1} z_i^{\pm 1}) = \prod_{1 \leq i \leq n} \theta_{q^{1/2}}(q^{1/4} v z_i^{\pm 1}) \quad (5.49)$$

on the left can be replaced by an arbitrary BC_n $q^{1/2}$ -theta function f of degree 1, in which case the extra factor on the right is

$$f(\pm \sqrt{-z_1}, \dots, \pm \sqrt{-z_{n/2}}), \quad \text{or} \quad f(\pm \sqrt{-z_1}, \dots, \pm \sqrt{-z_{(n-1)/2}}, \sqrt{-1}), \quad (5.50)$$

as appropriate. The factor on the right vanishes iff

$$f(-z_1, \dots, -z_n) = -f(z_1, \dots, z_n). \quad (5.51)$$

Remark 2. In fact, algebraically speaking, there is another modular transformation, since this conjecture depends on an ordered pair of 2-torsion points (namely -1 and $p^{1/2}$), so there is a “vanishing” conjecture associated to the pair $(\pm p^{1/2})$. However, this conjecture does not appear amenable to extension to a full integral.

The corresponding “vanishing” result states that the integral

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots; p^{1/4} t_0; p^{-1/4} t_0, p^{1/4} t_1, p^{-1/4} t_1; p^{1/4} u_0, p^{-1/4} u_0; q, t; p) \rangle_{t_0, t_1, u_0, 1, t^{1/2}, q^{1/2} t^{1/2}; t; p^{1/2}, q}^{(n/2)} \quad (5.52)$$

or

$$\langle \tilde{\mathcal{R}}_{\lambda}^{(n)}(\dots, p^{1/4} z_i^{\pm 1}, \dots, p^{1/4}; p^{1/4} t_0; p^{-1/4} t_0, p^{1/4} t_1, p^{-1/4} t_1; p^{1/4} u_0, p^{-1/4} u_0; q, t; p) \rangle_{t_0, t_1, u_0, t, t^{1/2}, q^{1/2} t^{1/2}; t; p^{1/2}, q}^{((n-1)/2)} \quad (5.53)$$

as appropriate, evaluates to

$$\frac{\Delta_{\lambda}(-1/u_0|t^{n/2}, t^{(n-1)/2}t_0, \pm(t^{n-1}t_0u_0)^{-1/2}; t^{1/2}; p^{1/2}, q^{1/2})}{\Delta_{\lambda}(1/u_0^2|t^n, t^{n-1}t_0^2, 1/t^{n-1}t_0u_0, 1/t^{n-1}t_0u_0; t; p, q)}. \quad (5.54)$$

Again, for $n = 1$, this is a known quadratic evaluation [16].

The conjecture obtained from Conjecture 11 in the corresponding way is the same, except with p and q swapped. (The fact that this changes a sum of two integrals to a single integral should not be a concern, since after all Conjectures 10 and 11 are each other's modular transforms.) We thus have only one more transform to consider, namely that obtained from Conjecture 9.

Conjecture 13. $(t^{-1/2}, -1)$ Subject to the balancing conditions $t^{n-1}t_0t_1t_2u_0 = pqt^{1/2}$, $v_0v_1 = p^{1/2}q^{1/2}/t^{1/2}$, one has

$$\begin{aligned} & \int \mathcal{R}_{\lambda}^{*(n)}(; t^{-1/4}t_0, t^{-1/4}u_0; t; p, q) \\ & \quad \Delta^{(n)}(; t^{-1/4}t_0, t^{-1/4}t_1, t^{-1/4}t_2, t^{-1/4}u_0, t^{1/4}, p^{1/2}t^{1/4}, q^{1/2}t^{1/4}, p^{1/2}q^{1/2}t^{1/4}, -t^{1/4}v_0^2, -t^{1/4}v_1^2; t^{1/2}; p, q) \\ &= \frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t^{-1/2}t_0t_1; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1; t; p, q)} \prod_{0 \leq i < n} \prod_{0 \leq r < s < 3} \frac{\Gamma_{p,q}(t^{i-1/2}t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s)} \\ & \quad \int \mathcal{R}_{\lambda}^{*(n)}(\dots, z_i^2, \dots; -t_0, -u_0; t; p, q) \Delta^{(n)}(; \pm\sqrt{-t_0}, \pm\sqrt{-t_1}, \pm\sqrt{-t_2}, \pm\sqrt{-u_0}, v_0, v_1; t^{1/2}; p^{1/2}, q^{1/2}). \end{aligned} \quad (5.55)$$

Proposition 5.9. Conjecture 13 holds for $n = 1$.

Proof. Exchange order of integration in the double integral

$$\int \int \Gamma_{p,q}(t^{1/4}y^{\pm 2}x^{\pm 1}) \frac{\prod_{0 \leq r < 4} \Gamma_{p,q}(u_r x^{\pm 1})}{\Gamma_{p,q}(x^{\pm 2})} \frac{dx}{2\pi\sqrt{-1}x} \frac{\Gamma_{p^{1/2}, q^{1/2}}(v_0 y^{\pm 1}, v_1 y^{\pm 1})}{\Gamma_{p^{1/2}, q^{1/2}}(y^{\pm 2})} \frac{dy}{2\pi\sqrt{-1}y}, \quad (5.56)$$

then use gamma function identities to express the integrands in the standard form. $\square$

The corresponding “vanishing” integral states that

$$\langle \tilde{R}_{\lambda}^{(n)}(; t^{-1/4}t_0; t^{1/4}t_0, t^{-1/4}t_1, t^{1/4}t_1; t^{-1/4}u_0, t^{1/4}u_0; t; p, q) \rangle_{t^{-1/4}t_0, t^{-1/4}t_1, t^{-1/4}u_0, t^{1/4}, p^{1/2}t^{1/4}, q^{1/2}t^{1/4}; t^{1/2}; p, q}^{(n)} \quad (5.57)$$

takes value (5.54).

We close with a combinatorial remark. The above seven conjectures (Conjectures 6, 7, 9, 10, 11, 12, 13), if we count both forms of those integrals not symmetric between p and q , give rise to twelve conjectures, one for each ordered pair (a, b) with $a \neq b \in \{-1, p^{1/2}, q^{1/2}, t^{-1/2}\}$. Furthermore, the three involutions “modular transform”, “swap p and q ”, and “dualize” act on the labels via their natural action on this set of square roots. Furthermore, in the conjecture associated to the pair (a, b) , the integrals are related by a factor

$$\frac{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1a; t; p, q)}{\Delta_{\lambda}^0(t^{n-1}t_0/u_0|t^{n-1}t_0t_1/b^2; t; p, q)} \prod_{\substack{0 \leq i < n \\ 0 \leq r < s < 3}} \frac{\Gamma_{p,q}(t^i t_r t_s)}{\Gamma_{p,q}(t^i t_r t_s / b^2)}, \quad (5.58)$$

with balancing condition $t^{n-1}t_0t_1t_2u_0 = pqb^2/a$, except that when $b = t^{-1/2}$, n must be replaced by $2n$. This pattern, together with corresponding patterns in the parameters of the interpolation functions, allows us to verify the consistency conditions for all of the cases at once, apart from checking that multiplying the interpolation functions by

$$\prod_{1 \leq i \leq n} \frac{\Gamma_{p,q}(a^{1/2}Qt_0z_i^{\pm 1}, a^{1/2}u_0z_i^{\pm 1}/Q)}{\Gamma_{p,q}(a^{1/2}t_0z_i^{\pm 1}, a^{1/2}u_0z_i^{\pm 1})}, \quad \text{or} \quad \prod_{1 \leq i \leq n} \frac{\Gamma_{p,q}(Qt_0z_i^{\pm 1}/b, u_0z_i^{\pm 1}/bQ)}{\Gamma_{p,q}(t_0z_i^{\pm 1}/b, u_0z_i^{\pm 1}/b)}, \quad (5.59)$$

as appropriate, before specializing, has the effect, after specializing, of shifting parameters in the corresponding integrand. Similarly, Conjectures 2, 3, and 4 correspond to four identities in natural bijection with the above four square roots. Unfortunately, there are enough quirks in the various cases to make it unclear how to formulate the conjectures in a more uniform manner.

References

- [1] H. Coskun and R. A. Gustafson. Well-poised Macdonald functions W_λ and Jackson coefficients ω_λ on BC_n . In *Jack, Hall-Littlewood and Macdonald polynomials*, volume 417 of *Contemp. Math.*, pages 127–155. Amer. Math. Soc., Providence, RI, 2006.
- [2] J. F. van Diejen and V. P. Spiridonov. An elliptic Macdonald-Morris conjecture and multiple modular hypergeometric sums. *Math. Res. Lett.*, 7(5-6):729–746, 2000.
- [3] J. F. van Diejen and V. P. Spiridonov. Elliptic Selberg integrals. *Internat. Math. Res. Notices*, (20):1083–1110, 2001.
- [4] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Univ. Press, Oxford, England, second edition, 1995.
- [5] A. Okounkov. BC -type interpolation Macdonald polynomials and binomial formula for Koornwinder polynomials. *Transform. Groups*, 3(2):181–207, 1998.
- [6] E. M. Rains. BC_n -symmetric polynomials. *Transform. Groups*, 10(1):63–132, 2005.
- [7] E. M. Rains. BC_n -symmetric abelian functions. *Duke Math. J.*, 135(1):99–180, 2006.
- [8] E. M. Rains. Transformations of elliptic hypergeometric integrals. *Ann. Math.*, to appear, 2009.
- [9] E. M. Rains and M. Vazirani. Vanishing integrals of Macdonald and Koornwinder polynomials. *Transform. Groups*, 12(4):725–759, 2007.
- [10] H. Rosengren. An elementary approach to $6j$ -symbols (classical, quantum, rational, trigonometric, and elliptic). *Ramanujan J.*, 13(1-3):131–166, 2007.
- [11] H. Rosengren and M. Schlosser. On Warnaar’s elliptic matrix inversion and Karlsson-Minton-type elliptic hypergeometric series. *J. Comput. Appl. Math.*, 178(1-2):377–391, 2005.

- [12] S. Sahi. Interpolation, integrality, and a generalization of Macdonald's polynomials. *Internat. Math. Res. Notices*, (10):457–471, 1996.
- [13] V. P. Spiridonov. On the elliptic beta function. *Uspekhi Mat. Nauk*, 56(1(337)):181–182, 2001.
- [14] V. P. Spiridonov. An elliptic incarnation of the Bailey chain. *Int. Math. Res. Not.*, (37):1945–1977, 2002.
- [15] S. O. Warnaar. Extensions of the well-poised and elliptic well-poised Bailey lemma. *Indag. Math. (N.S.)*, 14(3-4):571–588, 2003.
- [16] S. O. Warnaar. Summation formulae for elliptic hypergeometric series. *Proc. Amer. Math. Soc.*, 133(2):519–527 (electronic), 2005.